

Review

Open Access



A review of advanced head and neck osteoradionecrosis

Erica J. Mayland, Larissa Sweeny

Department of Otolaryngology - Head and Neck Surgery, Louisiana State University Health Science Center - New Orleans, New Orleans, LA 70809, USA.

Correspondence to: Dr. Larissa Sweeny, Department of Otolaryngology-Head and Neck Surgery, Louisiana State University Health Science Center - New Orleans, New Orleans, LA 70809, USA. E-mail: lswee1@lsuhsc.edu

How to cite this article: Mayland EJ, Sweeny L. A review of advanced head and neck osteoradionecrosis. *Plast Aesthet Res* 2021;8:62. <https://dx.doi.org/10.20517/2347-9264.2021.38>

Received: 20 Apr 2021 **First Decision:** 19 Jul 2021 **Revised:** 19 Jul 2021 **Accepted:** 22 Oct 2021 **Published:** 5 Dec 2021

Academic Editors: Matthew Spector, Marten Basta **Copy Editor:** Xi-Jun Chen **Production Editor:** Xi-Jun Chen

Abstract

Osteoradionecrosis (ORN) of the head and neck can be a devastating complication following radiation therapy. ORN is associated with pain, chronic infection, and non-healing wounds. Radiation fibrosis, chronic infection, fistula formation, and necrotic tissues can make treatment challenging. The following review article is a narrative on the management of advanced head and ORN.

Keywords: Head and neck reconstruction, free flap, osteoradionecrosis

INTRODUCTION

The majority of patients with advanced head and neck cancer will receive radiation as part of their treatment^[1]. For a subset of patients, radiation treatment has devastating side effects on the surrounding non-cancerous tissues. One of the most crippling outcomes is the development of osteoradionecrosis (ORN) in the head and neck region. These cases can range from mildly symptomatic areas of exposed bone to pathological fractures, often leaving patients with disabling symptoms. The current understanding of ORN pathophysiology suggests a progression of hypovascular-hypoxic-hypocellular tissue. Changes in the metabolic homeostasis following radiation eventually progress to a state of hypoxia and hypovascularity. This ultimately leads to tissue breakdown and a non-healing wound^[2].



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



The mean duration from completion of radiation therapy to the development of head and neck ORN is estimated at 22-47 months^[3]. Dental extractions are commonly found to be a precipitating factor, with some studies noting a recent dental extraction in 50%-60% of cases^[4,5]. The incidence of head and neck ORN throughout the literature ranges from 3%-15%^[3,6]. While ORN can occur in multiple head and neck subsites, the mandible is the most common location^[7]. The treatment of ORN is dependent on the severity of symptoms. The following narrative focuses on patients with advanced head and neck ORN with an emphasis on refractory cases requiring a free flap reconstruction.

CLINICAL PRESENTATION

Patients with advanced ORN commonly present with pain and exposed bone [Figure 1]. Patients with malignancy can also present with pain; therefore, an underlying malignancy should be ruled out prior to the start of ORN treatment. As ORN progresses, the bone may experience loss of density and strength, resulting in a pathologic fracture and/or orocutaneous fistula.

On panorex, early stages of ORN may present with findings of sclerotic bone or a poorly defined radiolucent lesion. Panorex may also show findings of cortication loss following dental extraction^[8]. Common computed tomography (CT) findings of ORN include cortical defect or lucency, disorganized bony architecture, intraosseous air, and ultimately pathologic fracture^[9]. When considering surgical intervention, CT imaging with 1mm cuts of the maxillofacial skeleton is recommended^[10].

NON-SURGICAL MANAGEMENT

ORN treatment typically begins conservatively, with free flap reconstruction being reserved for refractory and advanced cases^[11]. Conservative therapy for the early-stage disease includes optimizing oral hygiene, eliminating dental disease, and the use of systemic antibiotics^[12]. While often non-curative, these conservative interventions can provide symptomatic relief and slow progression. Antibiotics are commonly administered for acute infections or in the setting of chronically draining orocutaneous fistulas [Table 1].

Hyperbaric oxygen (HBO) therapy was introduced as a possible treatment for ORN in 1983 by Marx^[13]. HBO therapy works by increasing local tissue oxygen concentrations, thereby promoting tissue epithelialization and bone regeneration. HBO therapy is sometimes used prophylactically in patients requiring dental extractions after radiation to potentially prevent ORN development. However, a randomized controlled trial of previously radiated patients requiring dental extraction found HBO therapy before dental trauma did not prevent ORN complications^[14]. Currently, there are limited data demonstrating efficacy for the use of HBO therapy in the treatment of ORN [Table 2]^[15-17].

PENTOCLO is an antioxidant therapy that consists of pentoxifylline, tocopherol, and clodronate. Previous publications suggest improved wound healing when patients with ORN were administered PENTOCLO^[18,19]. Pentoxifylline is thought to improve microcirculation, while tocopherol (vitamin E) acts as an antioxidant. Clodronate is a first-generation bisphosphonate that reduces osteoclast activity and stimulates osteoblasts. A retrospective study found that patients who received pentoxifylline and tocopherol after radiation had a lower incidence of ORN^[20]. Dissard *et al.*^[21] also found that administration of the PENTOCLO regimen daily had a low side effect profile with high rates of symptom improvement when given antibiotics and steroids.

SURGICAL MANAGEMENT

ORN has significant quality of life (QOL) implications, including pain, infection, draining fistulas, and/or pathologic fractures. The decision to proceed with more invasive treatments is typically dictated by the

Table 1. Summary of literature evaluating the efficacy of antibiotic therapy in radiated patients receiving dental extraction or implantation

Ref.	Design	n	Treatment	Outcomes	Conclusion
Sandhu <i>et al.</i> ^[55] (2020)	Retrospective	50	Post dental extraction Amoxicillin 500 mg 3×/day for 14 days with chlorhexidine 2×/day in radiated patients	2% developed ORN at 18 months	ORN incidence was comparable with reported rates in the literature in radiated patients receiving post-extraction Abx
Al-Bazie <i>et al.</i> ^[56] (2016)	RCT	89	Ten days of Amoxicillin 500 mg q8h and chlorhexidine before extractions	No reported cases of ORN at a mean of 63 months	Perioperative Abx with antibiotic mouthwashes are effective in preventing ORN after extractions
Sultan <i>et al.</i> ^[15] (2017)	RCT	26	Abx alone vs. Abx + HBO prophylaxis in radiated patients receiving dental implants	1/13 from HBO + Abx developed ORN, 0/13 from Abx alone developed ORN No difference in implant survival	No difference in outcomes between Abx vs. Abx with HBO in radiated patients undergoing dental implants
Marx <i>et al.</i> ^[57] (1985)	RCT	37	Penicillin prior to dental extraction in radiated patients vs. HBO	ORN in 5.4% of HBO group and 29.9% of Abx group	Perioperative Abx did not decrease ORN incidence at 6 months following dental extraction

RCT: Randomized control trial; Abx: antibiotics; ORN: osteoradionecrosis; HBO: hyperbaric oxygen.

Table 2. Summary of literature evaluating the efficacy of hyperbaric oxygen therapy in patients with osteonecrosis

Ref.	Design	n	Treatment	Outcomes	Conclusion
Shaw <i>et al.</i> ^[14] (2019)	RCT	144	HBO vs. no HBO in radiated patients requiring dental extraction/implantation	Incidence of ORN at 6 months was 6.4% (HBO) vs. 5.7% (control)	HBO for dental extraction/implantation is unnecessary
Bennett <i>et al.</i> ^[58] (2016)	Cochrane Review	753	HBO vs. no HBO in patients with non-healing wounds	Improved healing of radiated sockets after extractions in HBO group	Suggest improved healing following HBO treatment in radiated sockets after extractions
Teguh <i>et al.</i> ^[59] (2009)	RCT	19	Thirty sessions of HBO after completing head and neck RT vs. no HBO	Higher QOL scores in HBO group (swallow, saliva, and pain)	Head and neck patients receiving HBO after RT had higher QOL scores
D'Souza <i>et al.</i> ^[60] (2007)	Retrospective	23	HBO vs. no HBO for treatment of ORN	12.5% cure in HBO group, 86% cure rate in non-HBO group	Small sample size but minimal benefit from HBO in the treatment of ORN
Bessereau <i>et al.</i> ^[61] (2010)	RCT	68	HBO vs. no HBO for treatment of ORN	None	Terminated early due to worse outcome in HBO arm

RCT: Randomized control trial; ORN: osteoradionecrosis; HBO: hyperbaric oxygen; QOL: quality of life.

patient’s symptomology and the subsequent impact on QOL. Conservative surgical measures can include debridement, sequestrectomy, local tissue rearrangement, and marginal mandibulectomy. If conservative medical and surgical management fails to provide resolution or symptom relief, more invasive surgical interventions can be considered. Definitive surgery typically involves resection of the involved bone and soft tissues. The resulting defect often requires reconstruction with vascularized non-radiated tissue, typically in the form of a free flap^[22-26].

FREE FLAP DONOR SITES

When selecting a donor site for ORN reconstruction, one must consider the amount of soft tissue required and the length of bone needed for continuity.

Fibula

The fibula is the donor site used most commonly for free flap reconstruction of head and neck ORN. Advantages of the fibula free flap include a long segment of bone stock (22-25 cm), adequate pedicle length for tension-free anastomosis in the neck, and low donor site morbidity^[27]. The fibula is harvested as a long




Figure 1. Physical examination of patient presenting with ORN following oral cavity radiation. Intraoral bone exposure (arrows) is a common finding in ORN patients often presenting as pain and oral phase dysphagia. ORN: Osteoradionecrosis.

segment of vascularized bone which allows for multiple osteotomies if needed to restore the natural contouring of the mandible or midface. The fibula has a high volume of cortical bone to bear the forces of mastication. Additionally, the bicortical bone of the fibula allows for dental implantation. The associated fasciocutaneous paddle has reliable perforators and can be harvested with the bone if coverage is needed for a mucosal or soft tissue defect. However, patients with severe peripheral vascular disease or those lacking three-vessel (anterior tibial, posterior tibial, peroneal) arterial runoff of the lower extremity may not be candidates for fibula reconstruction.

Scapula

In cases where the fibula is contraindicated or a large component of soft tissue is required, the scapula free flap is an ideal donor site. The lateral border of the scapula can reliably provide 10 cm of bone. Compared to the fibula, the scapula may have a thinner bone. The subscapular system often has a shorter pedicle length. However, the scapula tip is supplied by the angular artery, which originates from the thoracodorsal system. It has been reported that the angular artery is able to supply up to 10 cm of the lateral scapula border^[28]. As a result, incorporation of the angular artery allows for increased pedicle length. The subscapular system can be extremely versatile, providing skin, muscle, and bone in a multitude of combinations. The latissimus can be incorporated and utilized for intraoral or external plate coverage. An additional chimeric component of rib and serratus muscle can be harvested with the serratus branch of the thoracodorsal vessels. The versatility of soft tissue combinations and bulk makes the scapular flap ideal for complex soft tissue reconstructions requiring a large volume of soft tissue and a shorter linear segment of bone^[29].

Dental implantation is also possible with scapula bone, and success rates are similar to the fibula^[28]. In addition, the scapula may be advantageous in elderly patients who have significant peripheral vascular disease. Additionally, patients tend to ambulate sooner, possibly reducing postoperative complications^[30]. The main disadvantages of the scapula are the shorter pedicle length, longer operative time, and intraoperative patient positioning.

Osteocutaneous radial forearm

The osteocutaneous radial forearm free flap (OCRFFF) is another less commonly used option to restore bony continuity. Similar to the scapula, the OCRFFF provides 10-12 cm of bone. However, the entire

diameter of the radius bone cannot be harvested, limiting it to a single cortex following the harvest of 40%-60% of the radius circumference. Subsequently, there is insufficient bone stock to support dental implantation and necessitates prophylactic plating the radius to prevent fracture^[31,32]. In addition, the fasciocutaneous tissue volume available is often less than the soft tissue available with the fibula.

Fasciocutaneous and myocutaneous only donor sites

In patients who do not require a vascularized segment of bone, a radial forearm or anterolateral thigh flap may be utilized. These flaps are applicable in ORN of the anterior skull base or lateral temporal bone for soft tissue coverage^[33]. The radial forearm free flap is a fasciocutaneous flap that is thin and pliable, with reliable pedicle length and high rates of flap survival. However, this flap contains limited subcutaneous adipose in the majority of patients and lacks a muscle component making it inadequate for defects with large volume loss. Conversely, the anterolateral thigh free flap (ALT) may be composed of a combination of skin, adipose, fascia, or muscle. Adipose allows for improved volume retention as it is less subject to atrophy and contracture over time. However, depending on the thickness of the ALT subcutaneous adipose, it may provide more bulk than desired.

OPERATIVE CONSIDERATIONS

When reconstructing with an osseous donor site, consideration of bone segment length is important. For the periosteum to provide sufficient blood supply, it is recommended that each bone segment measure at least 2 cm in length. The ability to create multiple osteotomies can improve facial contouring. However, it also reduces the length of the pedicle. This can create additional challenges in vessel depleted, radiated necks where contralateral vessels or vessels in the base of the neck (transverse cervical, dorsal scapular, or internal mammary) may necessitate a longer pedicle length.

VIRTUAL SURGICAL PLANNING

Virtual surgical planning (VSP) can be utilized to aid in flap design preoperatively. VSP allows for the creation of patient-specific cutting guides and plates, which may increase free flap accuracy and reduce operative times^[34,35]. The use of VSP is thought to improve bone-to-bone contact, reducing rates of malunion or nonunion and the subsequent sequelae^[36,37]. The VSP software also provides information regarding the thickness of bone to aid in selecting sites for screw placement, ensuring adequate bone thickness, and reducing screw mobility and subsequent hardware extrusion^[36-38]. VSP can also be used to mirror the healthy bone allowing for more symmetric reconstruction that more accurately reflects the anatomical positioning prior to developing ORN. Additionally, VSP can aid in planning the ideal orientation of the pedicle^[39].

TECHNICAL CONSIDERATIONS

Flap geometry and vessel selection can be quite challenging in patients with ORN. Tissue damage and fibrosis from prior radiation therapy and chronic infection can complicate dissection of recipient vasculature and impact the vessel wall integrity and caliber. A previous publication found that over half of free flap reconstructions for ORN required the use of the contralateral neck vessels^[17]. Similarly, we have found improved free flap survival rates when using contralateral neck vessels^[40].

Reconstruction bar thickness varies among surgeons, ranging from 1 to 3 mm and averages 2.1 mm. An example of a 2.0 mm reconstruction bar is shown in [Figure 2A](#). Recent data suggests that reconstruction bar thickness may correlate with overlying soft tissue loss and hardware exposure rates, with a higher incidence of complications as reconstruction bar thickness increases^[40].




Figure 2. (A) Fibula free flap reconstruction of osteonecrosis defect extending from mandibular symphysis to ramus. (B) Example of the fasciocutaneous paddle of the fibula being to provide intraoral coverage for a mucosal defect.

Another variation in surgical design includes orientation and placement of the soft tissue component. Orientation of the fasciocutaneous paddle, in particular when a fibula donor site is used, can either be medial to the reconstructed bone segment or lateral to the reconstructed segment. When the soft tissue is oriented lateral to the hardware, it may provide an additional layer of soft tissue coverage over the reconstruction hardware. An example of a fasciocutaneous paddle that was brought medially into the floor of the mouth is depicted in [Figure 2B](#). Interestingly, the medial orientation of the fasciocutaneous paddle was found to result in a lower incidence of postoperative bone exposure and required fewer postoperative tissue debridement and local flaps procedures^[40]. The authors hypothesize that the medial orientation for the fasciocutaneous paddle reduces tension on the perforating vessels and provides protection from vascular insufficiency to the soft tissues^[40].

Dental implantation can significantly improve a patient's quality of life. Dental implantation success in ORN reconstruction has been cited around 95%^[41]. The literature cites similar complications rates with primary *vs.* secondary implantation in the setting of ORN. Secondary implantation has been found to correlate with higher fixed costs^[41], while primary implantation leads to a faster return of oral intake^[41]. Implant success rates were found to be similar for fibula *vs.* scapula free flaps^[42].

QUALITY OF LIFE

Pain is thought to be a large contributor to poor quality of life for patients with advanced ORN^[23,43]. Exposed nerve endings are subjected to infection and inflammation within a non-healing ORN wound. Free flaps are advantageous in their ability to provide coverage of exposed nerves and improve the blood flow to the area. One prospective study found a consistent reduction in pain-related domains following surgical resection and immediate free flap reconstruction of ORN, leading the authors to conclude that free flap reconstruction of advanced ORN improves QOL^[43].

In addition to pain, other QOL variables of concern for patients included speech, chewing, swallowing, and appearance^[24,44-46]. The University of Washington Quality of Life (UW-QOL) survey is commonly utilized for the assessment of QOL following head and neck reconstruction with a free flap^[23,24,43,46-48]. It was previously found that UW-QOL domain scores following free flap reconstruction for advanced ORN were higher for those patients who received dental implants, did not have a history of prior head and neck surgery, and did not develop a cancer recurrence^[22]. In contrast, Sweeny *et al.*^[22] found UW-QOL domain scores were not impacted by ORN recurrence, anatomic subsite of the ORN, or donor tissue used for the reconstruction. A summary of QOL outcomes can be found in [Table 3](#).

Table 3. Summary of literature evaluating long-term quality of life outcomes in patients following microvascular reconstruction of osteonecrosis defects

Ref.	Design	n	Treatment	Outcomes	Conclusion
Sweeny et al. ^[22] (2021)	Retrospective	137	UW-QOL survey in patients following free flap reconstruction for ORN	45% reported no pain, 28% no swallowing abnormalities, 93% no speech difficulty	Data suggests a good return of function and QOL following surgery
Lofstrand et al. ^[26] (2018)	Retrospective	41	SF-36, EORTC QLQ-C30, and QLQ-H&N35 questionnaires in cancer vs. ORN patients	ORN group had lower scores in swallowing and social eating compared to cancer, but general QOL did not differ from the reference population	Cancer and ORN patients have similar QOL following reconstruction with the exception of swallowing/social eating
Jacobson et al. ^[45] (2013)	Retrospective	30	PSS, SHI, QLQ-H&N35, and EAT-10 surveys in ORN after reconstruction	89% had abnormal EAT-10 and SHI scores following reconstruction, indicating abnormal speech and swallow	Many patients remain unhappy with speech and swallowing outcomes following reconstruction
Wang et al. ^[24] (2009)	Retrospective	15	UW-QOL in ORN after reconstruction	70% improved health related QOL after reconstruction, lowest scores in speech/swallow/saliva	Best scores in pain, but patients still have QOL issues with speech, swallow, and saliva

UW-QOL: University of Washington Quality of Life; SF-36: Short Form Health Survey; EORTC QLQ-C30: European Organization for Research and Treatment of Cancer quality of life questionnaire; QLQ-H&N35: European Organization for Research and Treatment of Cancer Quality of Life Questionnaire Head and Neck Module; PSS: Performance Status Scale; SHI: Speech Handicap Index; EAT-10: Eating Assessment Tool.

An important component of quality of life includes nutritional status. At 3 months following free flap reconstruction for advanced ORN, the rates of feeding tube dependence ranged from 13%-16%^[22,44]. When compared to their preoperative nutritional status, 47% of patients were tolerating a regular diet at 5 years following free flap reconstruction, and 31% had improvement in their diet status following free flap reconstructive surgery^[22]. This data suggest that for a subset of patients, free flap reconstruction can lead to an improvement in diet function.

SURGICAL COMPLICATIONS

Complication rates following free flap reconstruction of head and neck ORN are cited between 30%-60%^[16,26,44,46,49,50]. While most institutions cite free flap survival rates following head and neck reconstruction as 95% or greater, free flap survival rates following ORN reconstruction are cited at 89-93%^[26,46,50,51]. Additionally, it was found that patients undergoing free flap reconstruction for ORN had a higher incidence of late complications compared to patients undergoing free flap reconstruction for malignancy^[3]. A retrospective study of 277 patients found that 24% of patients developed a postoperative fistula, 16% developed exposed bone, and 20% developed plate extrusion following free flap reconstruction for ORN^[40]. These complications are attributed to poor tissue quality (radiation fibrosis, decreased perfusion) and a chronic inflammatory reaction in response to persistent infection and saliva exposure. Although uncommon, these complications can result in increased patient morbidity and healthcare costs^[50,52,53].

ORN RECURRENCE

ORN recurrence following free flap reconstruction is cited at 10%-14%, with a median time to onset of 11 months^[17,22]. Poor wound healing and failure of osseointegration postoperatively were found to correlate with higher rates of ORN recurrence^[22]. While the donor site selected for free flap reconstruction did not impact the recurrence of ORN^[54]. To avoid ORN recurrence at the surgical site, every attempt should be made to resect necrotic non-viable bone. The periosteum of the remaining bone should be inspected to ensure it is viable. The periosteum is a dense fibrous membrane with a rich vascular supply that

envelopes the outer cortex of the bone and provides the superficial cortex with nutrients. The cortex should also be evaluated for viability. The cortex of healthy bone is dense, typically white in color, and will have visible bright red bleeding from trans-cortical capillaries. In addition, the bone marrow should be evaluated for viability. Healthy bone marrow is able to maintain its trabecular structure and will bleed bright red. Any bone which does not appear viable should be removed until viable bone is confirmed.

CONCLUSIONS

ORN is one of the most crippling complications following radiation for head and neck malignancies. The development of ORN has a significant impact on quality of life, leaving patients with disabling pain and chronic wounds. Free flap reconstruction is an integral part of the definitive management of advanced ORN. Although often effective at treating ORN, there remains a risk for postoperative wound complications and functional decline. Informed discussions with patients regarding expectations and anticipated outcomes and careful preoperative planning are essential in treating these complex cases.

DECLARATIONS

Authors' contributions

Made substantial contributions to conception and design of the study, analysis and interpretation, composition of manuscript, and final approval of manuscript: Sweeny L, Mayland EJ

Availability of data and materials

Not applicable.

Financial support and sponsorship

None.

Conflicts of interest

Both authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2021.

REFERENCES

1. Trotti A, Bellm LA, Epstein JB, et al. Mucositis incidence, severity and associated outcomes in patients with head and neck cancer receiving radiotherapy with or without chemotherapy: a systematic literature review. *Radiother Oncol* 2003;66:253-62. [DOI PubMed](#)
2. Marx RE. Osteoradionecrosis: a new concept of its pathophysiology. *J Oral Maxillofac Surg* 1983;41:283-8. [DOI PubMed](#)
3. Danielsson D, Gahm C, Haghdoost S, Munck-Wikland E, Halle M. Osteoradionecrosis, an increasing indication for microvascular head and neck reconstruction. *Int J Oral Maxillofac Surg* 2020;49:1-6. [DOI PubMed](#)
4. Thorn JJ, Hansen HS, Specht L, Bastholt L. Osteoradionecrosis of the jaws: clinical characteristics and relation to the field of irradiation. *J Oral Maxillofac Surg* 2000;58:1088-93; discussion 1093-5. [DOI PubMed](#)
5. Aarup-Kristensen S, Hansen CR, Forner L, Brink C, Eriksen JG, Johansen J. Osteoradionecrosis of the mandible after radiotherapy for head and neck cancer: risk factors and dose-volume correlations. *Acta Oncol* 2019;58:1373-7. [DOI PubMed](#)
6. Moon DH, Moon SH, Wang K, et al. Incidence of, and risk factors for, mandibular osteoradionecrosis in patients with oral cavity and oropharynx cancers. *Oral Oncol* 2017;72:98-103. [DOI PubMed](#)
7. Reuther T, Schuster T, Mende U, Kübler A. Osteoradionecrosis of the jaws as a side effect of radiotherapy of head and neck tumour patients--a report of a thirty year retrospective review. *Int J Oral Maxillofac Surg* 2003;32:289-95. [DOI PubMed](#)
8. AlDhalaan NA, BaQais A, Al-Omar A. Medication-related osteonecrosis of the jaw: a review. *Cureus* 2020;12:e6944. [DOI PubMed](#)

PMC

9. Jereczek-Fossa BA, Orecchia R. Radiotherapy-induced mandibular bone complications. *Cancer Treat Rev* 2002;28:65-74. DOI PubMed
10. Ren W, Gao L, Li S, et al. Virtual Planning and 3D printing modeling for mandibular reconstruction with fibula free flap. *Med Oral Patol Oral Cir Bucal* 2018;23:e359-66. DOI PubMed PMC
11. Oh HK, Chambers MS, Martin JW, Lim HJ, Park HJ. Osteoradionecrosis of the mandible: treatment outcomes and factors influencing the progress of osteoradionecrosis. *J Oral Maxillofac Surg* 2009;67:1378-86. DOI PubMed
12. Khan AA, Morrison A, Hanley DA, et al; International Task Force on Osteonecrosis of the Jaw. Diagnosis and management of osteonecrosis of the jaw: a systematic review and international consensus. *J Bone Miner Res* 2015;30:3-23. DOI PubMed
13. Marx RE. A new concept in the treatment of osteoradionecrosis. *J Oral Maxillofac Surg* 1983;41:351-7. DOI PubMed
14. Shaw RJ, Butterworth CJ, Silcocks P, et al. HOPON (hyperbaric oxygen for the prevention of osteoradionecrosis): a randomized controlled trial of hyperbaric oxygen to prevent osteoradionecrosis of the irradiated mandible after dentoalveolar surgery. *Int J Radiat Oncol Biol Phys* 2019;104:530-9. DOI PubMed
15. Sultan A, Hanna GJ, Margalit DN, et al. The use of hyperbaric oxygen for the prevention and management of osteoradionecrosis of the jaw: a Dana-Farber/Brigham and Women's Cancer Center Multidisciplinary Guideline. *Oncologist* 2017;22:343-50. DOI PubMed PMC
16. Hirsch DL, Bell RB, Dierks EJ, Potter JK, Potter BE. Analysis of microvascular free flaps for reconstruction of advanced mandibular osteoradionecrosis: a retrospective cohort study. *J Oral Maxillofac Surg* 2008;66:2545-56. DOI PubMed
17. Alam DS, Nuara M, Christian J. Analysis of outcomes of vascularized flap reconstruction in patients with advanced mandibular osteoradionecrosis. *Otolaryngol Head Neck Surg* 2009;141:196-201. DOI PubMed
18. Delanian S, Depondt J, Lefaix JL. Major healing of refractory mandible osteoradionecrosis after treatment combining pentoxifylline and tocopherol: a phase II trial. *Head Neck* 2005;27:114-23. DOI PubMed
19. Delanian S, Chatel C, Porcher R, Depondt J, Lefaix JL. Complete restoration of refractory mandibular osteoradionecrosis by prolonged treatment with a pentoxifylline-tocopherol-clodronate combination (PENTOCLO): a phase II trial. *Int J Radiat Oncol Biol Phys* 2011;80:832-9. DOI PubMed
20. Patel V, Gadiwalla Y, Sassoon I, Sproat C, Kwok J, McGurk M. Prophylactic use of pentoxifylline and tocopherol in patients who require dental extractions after radiotherapy for cancer of the head and neck. *Br J Oral Maxillofac Surg* 2016;54:547-50. DOI PubMed
21. Dissard A, P Dang N, Barthelemy I, et al. Efficacy of pentoxifylline-tocopherol-clodronate in mandibular osteoradionecrosis. *Laryngoscope* 2020;130:E559-66. DOI PubMed
22. Sweeny L, Mayland E, Swendseid BP, et al. Microvascular reconstruction of osteonecrosis: assessment of long-term quality of life. *Otolaryngol Head Neck Surg* 2021. DOI PubMed
23. Rogers SN, D'Souza JJ, Lowe D, Kanatas A. Longitudinal evaluation of health-related quality of life after osteoradionecrosis of the mandible. *Br J Oral Maxillofac Surg* 2015;53:854-7. DOI PubMed
24. Wang L, Su YX, Liao GQ. Quality of life in osteoradionecrosis patients after mandible primary reconstruction with free fibula flap. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;108:162-8. DOI PubMed
25. Shaha AR, Cordeiro PG, Hidalgo DA, et al. Resection and immediate microvascular reconstruction in the management of osteoradionecrosis of the mandible. *Head Neck* 1997;19:406-11. DOI PubMed
26. Lofstrand J, Nyberg M, Karlsson T, et al. Quality of life after free fibula flap reconstruction of segmental mandibular defects. *J Reconstr Microsurg* 2018;34:108-20. DOI PubMed
27. Kim JW, Hwang JH, Ahn KM. Fibular flap for mandible reconstruction in osteoradionecrosis of the jaw: selection criteria of fibula flap. *Maxillofac Plast Reconstr Surg* 2016;38:46. DOI PubMed PMC
28. Blumberg JM, Walker P, Johnson S, et al. Mandibular reconstruction with the scapula tip free flap. *Head Neck* 2019;41:2353-8. DOI PubMed
29. Yeh DH, Lee DJ, Sahovaler A, et al. Shouldering the load of mandible reconstruction: 81 cases of oromandibular reconstruction with the scapular tip free flap. *Head Neck* 2019;41:30-6. DOI PubMed
30. Dowthwaite SA, Theurer J, Belzile M, et al. Comparison of fibular and scapular osseous free flaps for oromandibular reconstruction: a patient-centered approach to flap selection. *JAMA Otolaryngol Head Neck Surg* 2013;139:285-92. DOI PubMed
31. Bigcas JLM, Bond J. Osteocutaneous radial forearm flap. Treasure Island (FL): StatPearls Publishing; 2021.
32. Kim JH, Rosenthal EL, Ellis T, Wax MK. Radial forearm osteocutaneous free flap in maxillofacial and oromandibular reconstructions. *Laryngoscope* 2005;115:1697-701. DOI PubMed
33. Chapchay K, Weinberger J, Eliashar R, Adler N. Anterior skull base reconstruction following ablative surgery for osteoradionecrosis: case report and review of literature. *Ann Otol Rhinol Laryngol* 2019;128:1134-40. DOI PubMed
34. Sharaf B, Levine JP, Hirsch DL, Bastidas JA, Schiff BA, Garfein ES. Importance of computer-aided design and manufacturing technology in the multidisciplinary approach to head and neck reconstruction. *J Craniofac Surg* 2010;21:1277-80. DOI PubMed
35. Largo RD, Garvey PB. Updates in head and neck reconstruction. *Plast Reconstr Surg* 2018;141:271e-85e. DOI PubMed
36. Avraham T, Franco P, Brecht LE, et al. Functional outcomes of virtually planned free fibula flap reconstruction of the mandible. *Plast Reconstr Surg* 2014;134:628e-34e. DOI PubMed
37. Chang EI. Long-term operative outcomes of preoperative computed tomography-guided virtual surgical planning for osteocutaneous free flap mandible reconstruction. *Plast Reconstr Surg* 2016;138:559e-60e. DOI PubMed
38. Patel H, Saadat N, Ho AS, Mallen-St Clair J. Virtual surgical planning for bisphosphonate-related osteonecrosis of the jaw: a valuable

- application in advanced cases. *Cureus* 2020;12:e9696. DOI PubMed PMC
39. Vieira L, Isacson D, Dimovska EOF, Rodriguez-Lorenzo A. Four lessons learned from complications in head and neck microvascular reconstructions and prevention strategies. *Plast Reconstr Surg Glob Open* 2021;9:e3329. DOI PubMed PMC
 40. Mayland EJ SB, Curry JM, Wax MK, et al. Management and and outcomes of advanced head and neck osteonecrosis. Forthcoming 2021.
 41. Menapace DC, Van Abel KM, Jackson RS, Moore EJ. Primary vs secondary endosseous implantation after fibular free tissue reconstruction of the mandible for osteoradionecrosis. *JAMA Facial Plast Surg* 2018;20:401-8. DOI PubMed PMC
 42. Wilkman T, Husso A, Lassus P. Clinical comparison of scapular, fibular, and iliac crest osseal free flaps in maxillofacial reconstructions. *Scand J Surg* 2019;108:76-82. DOI PubMed
 43. Danielsson D, Munck-Wikland E, Hagel E, Halle M. Quality of life after microvascular mandibular reconstruction for osteoradionecrosis-a prospective study. *Head Neck* 2019;41:2225-30. DOI PubMed
 44. Baumann DP, Yu P, Hanasono MM, Skoracki RJ. Free flap reconstruction of osteoradionecrosis of the mandible: a 10-year review and defect classification. *Head Neck* 2011;33:800-7. DOI PubMed
 45. Jacobson AS, Zevallos J, Smith M, et al. Quality of life after management of advanced osteoradionecrosis of the mandible. *Int J Oral Maxillofac Surg* 2013;42:1121-8. DOI PubMed
 46. Chang EI, Leon P, Hoffman WY, Schmidt BL. Quality of life for patients requiring surgical resection and reconstruction for mandibular osteoradionecrosis: 10-year experience at the University of California San Francisco. *Head Neck* 2012;34:207-12. DOI PubMed
 47. Rogers SN, Gwanne S, Lowe D, Humphris G, Yueh B, Weymuller EA Jr. The addition of mood and anxiety domains to the University of Washington quality of life scale. *Head Neck* 2002;24:521-9. DOI PubMed
 48. Derek Lowe SR. University of Washington Quality of Life Questionnaire (UW-QOL v4) Guidance for scoring and presentation. Available from: https://www.entnet.org/wp-content/uploads/files/uw_qol_r_v4.pdf [Last accessed on 1 Nov 2021].
 49. Sweeny L, Rosenthal EL, Light T, et al. Effect of overlapping operations on outcomes in microvascular reconstructions of the head and neck. *Otolaryngol Head Neck Surg* 2017;156:627-35. DOI PubMed
 50. Cannady SB, Dean N, Kroeker A, Albert TA, Rosenthal EL, Wax MK. Free flap reconstruction for osteoradionecrosis of the jaws--outcomes and predictive factors for success. *Head Neck* 2011;33:424-8. DOI PubMed
 51. Lee M, Chin RY, Eslick GD, Sriharan N, Paramaesvaran S. Outcomes of microvascular free flap reconstruction for mandibular osteoradionecrosis: a systematic review. *J Craniomaxillofac Surg* 2015;43:2026-33. DOI PubMed
 52. Park L, Lilic N, Addison B, Patel R. Cost analysis of osteoradionecrosis. *J Laryngol Otol* 2017;131:303-8. DOI PubMed
 53. Kelishadi SS, St-Hilaire H, Rodriguez ED. Is simultaneous surgical management of advanced craniofacial osteoradionecrosis cost-effective? *Plast Reconstr Surg* 2009;123:1010-7. DOI PubMed
 54. Li H, Tan MDM, Alexander S, Grinsell D, Ramakrishnan A. Comparative osteoradionecrosis rates in bony reconstructions for head and neck malignancy. *J Plast Reconstr Aesthet Surg* 2019;72:1478-83. DOI PubMed
 55. Sandhu S, Salous MH, Sankar V, Margalit DN, Villa A. Osteonecrosis of the jaw and dental extractions: a single-center experience. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2020;130:515-21. DOI PubMed
 56. Al-Bazie SA, Bahatheq M, Al-Ghazi M, Al-Rajhi N, Ramalingam S. Antibiotic protocol for the prevention of osteoradionecrosis following dental extractions in irradiated head and neck cancer patients: a 10 years prospective study. *J Cancer Res Ther* 2016;12:565-70. DOI PubMed
 57. Marx RE, Johnson RP, Kline SN. Prevention of osteoradionecrosis: a randomized prospective clinical trial of hyperbaric oxygen versus penicillin. *J Am Dent Assoc* 1985;111:49-54. DOI PubMed
 58. Bennett MH, Feldmeier J, Hampson NB, Smee R, Milross C. Hyperbaric oxygen therapy for late radiation tissue injury. *Cochrane Database Syst Rev* 2016;4:CD005005. DOI PubMed PMC
 59. Teguh DN, Levendag PC, Noever I, et al. Early hyperbaric oxygen therapy for reducing radiotherapy side effects: early results of a randomized trial in oropharyngeal and nasopharyngeal cancer. *Int J Radiat Oncol Biol Phys* 2009;75:711-6. DOI PubMed
 60. D'Souza J, Goru J, Goru S, Brown J, Vaughan ED, Rogers SN. The influence of hyperbaric oxygen on the outcome of patients treated for osteoradionecrosis: 8 year study. *Int J Oral Maxillofac Surg* 2007;36:783-7. DOI PubMed
 61. Bessereau J, Annane D. Treatment of osteoradionecrosis of the jaw: the case against the use of hyperbaric oxygen. *J Oral Maxillofac Surg* 2010;68:1907-10. DOI PubMed