

Supplementary Table 1. Relationship between medical treatment and the incidence of olfactory and gustatory dysfunctions

Medication	All patients	Olfactory dysfunction	χ^2	P-value	Gustatory dysfunction	χ^2	P-value
	(n = 81)	(n = 11)			(n = 6)		
Classification of medication							
Antiviral	72(88.9)	10(90.9)	0.053	1.000*	6(100.0)	0.81	1.000*
γ - globulin biologics	36(44.4)	5(45.5)	0.005	1.000*	1(16.7)	2.025	0.219*
Chloroquine	26(32.1)	4(36.4)	0.106	0.739*	3(50.0)	0.953	0.380*
Glucocorticoids	32(39.5)	2(18.2)	2.422	0.186*	0(0.0)	4.232	0.076*
Thymosin	25(30.9)	5(45.5)	1.27	0.301*	4(66.7)	3.893	0.070*
Interferon	13(16.0)	3(27.3)	1.19	0.371*	3(50.0)	5.544	0.051
Antibiotics	57(70.4)	7(63.6)	0.277	0.724*	4(66.7)	0.043	1.000*
Combination therapy	78(96.3)	11(100.0)	0.49	1.000*	6(100.0)	0.249	1.000*

P-value denoted the comparison between different groups. *Fisher exact test.