

Supplementary Materials

Disease-specific signatures of circulating extracellular vesicles detected by the surface plasmon resonance imaging: a pilot study

Tatsuki Shibuta¹, Yukichi Takada¹, Shiori Nishinosono¹, Seiko Yasuda¹, Yasuhiro Ono², Yoshitaka Hirooka^{1,3,4}, Daisuke Irikura⁵, Kensuke Saito⁵, Tsukuru Umemura^{1,4}

¹Department of Medical Technology and Sciences, International University of Health and Welfare, Fukuoka 831-8501, Japan.

²Department of Diabetes and Metabolism, Kouhoukai Takagi Hospital, Fukuoka 831-0016, Japan.

³Hypertension and Heart Failure Center, Kouhoukai Takagi Hospital, Fukuoka 831-0016, Japan.

⁴Graduate School, International University of Health and Welfare, Fukuoka 831-8501, Japan.

⁵HORIBA, Ltd, Kyoto City, Kyoto 601-8510, Japan.

Correspondence to: Prof. Tatsuki Shibuta, Department of Medical Technology and Sciences, International University of Health and Welfare, 137-1 Enokizu, Okawa City, Fukuoka 831-8501, Japan. E-mail: tshibuta@iuhw.ac.jp; Dr. Tsukuru Umemura, Graduate School, International University of Health and Welfare, 137-1 Enokizu, Okawa City, Fukuoka 831-8501, Japan. E-mail: umemura@iuhw.ac.jp

Supplementary Table 1. Post hoc power analysis

EV markers	T2DM		n needed for 80% power	HT	
	Effect size	Power		Effect size	Power
CD20	5.44	1.00	-	5.76	1.00
DARC	3.85	1.00	-	0.83	0.84
CD81	2.02	1.00	-	2.06	1.00
AcvRIIa	4.43	1.00	-	5.61	1.00
CD9	5.93	1.00	-	1.72	1.00
EpoR	0.60	0.62	-	3.60	1.00
CD110	4.16	1.00	-	0.80	0.81
CD71	2.07	1.00	-	2.87	1.00
TfR2	2.59	1.00	-	3.75	1.00
CD44	1.03	0.97	-	0.71	0.71
MBM	0.95	0.94	-	2.75	1.00
CD4	0.16	0.09*	635	0.81	0.82
CD8	0.19	0.11*	420	0.94	0.91
CD25	0.27	0.17*	217	0.70	0.70
CD169	0.07	0.06*	3102	0.78	0.79

*If power is less than 0.6, the required sample size is calculated. T2DM: Type II diabetes mellitus; HT: hypertension; DARC: Duffy antigen receptor for chemokine; AcvRIIa: activin receptor type-2A; EpoR: erythropoietin receptor; TfR2: transferrin receptor 2; MBM: MAM lectin binding molecules.

CD44	<i>0.64</i>	-	-0.131	-	-	-	0.049	-0.243	-0.279	-0.012	0.154	-	0.050	-0.152
	7	0.13		0.08	0.12	0.17						0.051		
		3		1	6	5								
MBM	-	-	-0.109	0.08	0.15	0.29	0.167	-0.086	0.184	0.030	-0.177	0.007	<i>0.755</i>	-0.126
	0.07	0.24		2	1	3								
	8	7												

The values in the table indicate correlation coefficients (r). Bold and italicized numbers are those with higher correlation coefficients than others. T2DM: Type II diabetes mellitus; HT: hypertension; eGFR: estimated Glomerular Filtration Rate; sCre: serum Creatinine; U-Alb: Urine-Albumin; DARC: Duffy antigen receptor for chemokine; AcvRIIa: activin receptor type-2A; EpoR: erythropoietin receptor; Tfr2: transferrin receptor 2; MBM: MAM lectin binding molecules.

Supplementary Table 3. Logistic regression analysis results on laboratory data and SPRI intensities in T2DM and HT patients

Variable	T2DM		Nephropathy		HT	
	Retinopathy (n = 21)	<i>P</i>	OR (95% CI)	<i>P</i>	Moderate/High risk Group (n = 22)	<i>P</i>
Ages	0.996 (0.94 - 1.05)	0.873	1.022 (0.93 - 1.12)	0.637	1.010 (0.95 - 1.08)	0.763
Gender	0.516 (0.16 - 1.71)	0.279	1.656 (0.38 - 2.10)	0.101	2.179 (0.33 - 1.44)	0.419
eGFR	0.991 (0.97 - 1.02)	0.461	1.133 (1.03 - 1.32)	0.088		
sCre	1.263 (0.66 - 2.41)	0.479	0.946 (0.47 - 1.63)	0.097		
U-Alb	1.001 (0.99 - 1.01)	0.398	1.074 (1.02 - 1.13)	0.005*		
CD20	0.315 (0.09 - 1.09)	0.067	1.169 (0.41 - 2.59)	0.237		
CD44	0.002 (0.00 - 1.08)	0.053	0.809 (0.25 - 0.87)	0.414		
BNP					0.997 (0.95 - 1.05)	0.965
Nad					1.023 (1.01 - 1.04)	0.003*
AcvRIIa					0.070 (0.00 - 0.24)	0.063
MBM					0.036 (0.00 - 0.91)	0.022*

* $P < 0.05$. Bold numbers without asterisks are low P-values that did not achieve the 0.05 criterion. T2DM: Type II diabetes mellitus; HT: hypertension; OR: odds ratio; CI: confidence interval; eGFR: estimated Glomerular Filtration Rate; sCre: serum Creatinine; U-Alb: Urine-Albumin; BNP: brain natriuretic peptide; Nad: noradrenaline; AcvRIIa: activin receptor type-2A; MBM: MAM lectin binding molecules.