

Supplementary Materials

Painful nerve compression beyond the carpal tunnel: recognizing the lacertus syndrome

Yasmeen Al-Hashimi¹, Benjamin Ferembach², Vincent Martinel³, Elisabet Hagert^{4,5}

¹Faculty of Medicine, Weill Cornell Medicine - Qatar, Doha 24144, Qatar.

²Department of Orthopedics, NCT+ (Nouvelle Clinique Tours), Saint Cyr sur Loire 37540, France.

³Department of Orthopedics, Clinique Ormeau Pyrenées, Tarbes 65000, France.

⁴Department of Surgery, Aspetar Orthopedic and Sports Medicine Hospital, Doha 29222, Qatar.

⁵Department of Clinical Science and Education, Karolinska Institutet, Stockholm 11883, Sweden.

Correspondence to: Dr. Elisabet Hagert, Department of Surgery, Aspetar Orthopedic and Sports Medicine Hospital, Sports City Street, Aspire Zone, Doha 29222, Qatar. E-mail: elisabet.hagert@aspetar.com

ORCID: Elisabet Hagert (0000-0002-6667-2593)

Video 1. Intraoperative video (French, English subtitles) showing the weakness in FPL, FDP II, FCR, and pinch strength before lacertus release and the immediate return of power following sectioning of the lacertus fibrosus.