

Supplementary Materials

Prognostic value of lymphocyte subset levels in hepatocellular carcinoma following conventionally fractionated vs. stereotactic body radiotherapy

Si-Tong Wang[#], Yi-Xing Chen[#], Yu-Nan Gao, Ping Yang, Qian-Qian Zhao, Zhao-Chong Zeng, Yuan Zhuang

Department of Radiation Oncology, Zhongshan Hospital, Fudan University, Shanghai 200032, China.

[#]Authors contributed equally.

Correspondence to: Dr. Yuan Zhuang, Prof. Zhao-Chong Zeng, Department of Radiation Oncology, Zhongshan Hospital, Fudan University, 180 Fenglin Road, Xuhui, Shanghai 200032, China. E-mail: zhuang.yuan@zs-hospital.sh.cn; zeng.zhaochong@zs-hospital.sh.cn

Supplementary Table 1. The percentage of lymphocyte subpopulations decreased after radiotherapy

Variables	SBRT	CFRT	<i>P</i>-value
TPLC	50.7%	66.4%	0.004
CD4	46.5%	66.3%	0.003
CD8	50.4%	52.6%	0.21
NK	47.3%	71.8%	0.002
B	78.9%	90.5%	0.004

CFRT: Conventionally fractionated radiotherapy; SBRT: stereotactic body radiotherapy; TPLC: total peripheral lymphocyte count.