

Review

Open Access



Multi-layered control of PD-L1 expression in Epstein-Barr virus-associated gastric cancer

Christos N. Miliotis, Frank J. Slack

HMS Initiative for RNA Medicine, Department of Pathology, Cancer Center, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02115, USA.

Correspondence to: Dr. Frank J. Slack, Department of Pathology, Cancer Center, Beth Israel Deaconess Medical Center, Harvard Medical School, 3 Blackfan Circle, Boston, MA 02115, USA. E-mail: fslack@bidmc.harvard.edu

How to cite this article: Miliotis CN, Slack FJ. Multi-layered control of PD-L1 expression in Epstein-Barr virus-associated gastric cancer. *J Cancer Metastasis Treat* 2020;6:13. <http://dx.doi.org/10.20517/2394-4722.2020.12>

Received: 1 Feb 2020 **First Decision:** 7 Apr 2020 **Revised:** 23 Apr 2020 **Accepted:** 12 May 2020 **Published:** 23 May 2020

Science Editor: Pravin D. Potdar **Copy Editor:** Jing-Wen Zhang **Production Editor:** Jing Yu

Abstract

Gastric cancer (GC) is the fifth most common cancer worldwide. In approximately 10% of GC cases, cancer cells show ubiquitous and monoclonal Epstein-Barr virus (EBV) infection. A significant feature of EBV-associated GC (EBVaGC) is high lymphocytic infiltration and high expression of immune checkpoint proteins, including programmed death-ligand 1 (PD-L1). This highlights EBVaGC as a strong candidate for immune checkpoint blockade therapy. Indeed, several recent studies have shown that EBV positivity in GC correlates with positive response to programmed cell death protein 1 (PD-1)/PD-L1 blockade therapy. Understanding the mechanisms that control PD-L1 expression in EBVaGC can indicate new predictive biomarkers for immunotherapy, as well as therapeutic targets for combination therapy. Various mechanisms have been implicated in PD-L1 expression regulation, including structural variations, post-transcriptional control, oncogenic activation of intrinsic signaling pathways, and increased sensitivity to extrinsic signals. This review provides the most recent updates on the multi-layered control of PD-L1 expression in EBVaGC.

Keywords: Epstein-Barr virus, gastric cancer, immunotherapy, programmed death-ligand 1

INTRODUCTION

Gastric cancer (GC) is the fifth most common and third most deadly cancer worldwide^[1]. Due to the lack of early symptoms, patients are usually diagnosed with locally advanced or metastatic cancer, when



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Table 1. Immunological and molecular characteristics of GC molecular subtypes

Molecular subtype	EBV-positive	MSI	CIN	GS
Frequency (TCGA STAD) % ^[4]	9	22	50	20
TIL abundance ^[5,6]	High	High	Low	Low
PD-L1 expression ^[4-6]	High	High	Low	Low
Somatic mutations ^[4]	Standard rate of mutations (non-hypermutated) Common <i>PI3KCA</i> , <i>ARID1A</i> mutations	Hypermutated Recurrent mutations in <i>TP53</i> , <i>KRAS</i> , <i>ARID1A</i> , <i>PIK3CA</i> , <i>ERBB3</i> , <i>PTEN</i> and <i>HLA-B</i>	Non-hypermutated Common <i>TP53</i> mutations	Non-hypermutated Common <i>RHOA</i> , <i>CDH1</i> mutations
Other molecular characteristics ^[4]	Hypermethylation (EBV-CIMP, frequent <i>CDKN2A</i> silencing) Amplification in 9p24.1 chromosomal region (<i>PD-L1/PD-L2/JAK2</i>)	Hypermethylation (MSI-associated gastric CIMP, frequent <i>MLH1</i> silencing)	Gene amplifications, frequent in TKRs (<i>EGFR</i> , <i>VEGFA</i>) and deletions	

*“High” indicates that the corresponding molecular subtype is typically associated with high levels of TIL or PD-L1, respectively. “Low” indicates that the corresponding molecular subtype is not typically associated with high levels of TIL or PD-L1, respectively. MSI: microsatellite instability; CIN: chromosomal instability; GS: genomically stable; TCGA: the cancer genome atlas; STAD: stomach adenocarcinoma; TILs: tumor infiltrating lymphocytes; PD-L1: programmed death-ligand 1; PIK3CA: phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha; ARID1A: AT-rich interaction domain 1A; TP53: tumor protein p53; KRAS: Kirsten rat sarcoma viral oncogene homolog; ERBB3: Erb-B2 receptor tyrosine kinase 3; PTEN: phosphatase and tensin homolog; HLA-B: human leukocyte antigen B; RHOA: ras homolog family member A; CDH1: cadherin 1; CIMP: CpG island methylator phenotype; CDKN2A: cyclin dependent kinase inhibitor 2A; JAK2: Janus kinase 2; MLH1: MutL homolog 1; TKR: tyrosine kinase receptor; EGFR: epidermal growth factor receptor; VEGFA: vascular endothelial growth factor A; GC: gastric cancer

conventional lines of therapy are less effective. The 5-year survival rate for GC in the United States varies by stage and ranges from 68% in patients with localized cancer to 5% in patients with distant metastases^[2].

The majority of GCs (90%-95%) are adenocarcinomas^[3]. The Cancer Genome Atlas (TCGA) recently undertook a comprehensive molecular characterization of hundreds of gastric adenocarcinomas and proposed classification of GC into four molecular subtypes: Epstein-Barr virus (EBV)-positive, microsatellite instability (MSI), chromosomal instability (CIN), and genomically stable (GS) subgroups^[4]. EBV-positive tumors are characterized by DNA hypermethylation, frequent phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha (*PIK3CA*) mutations, and programmed cell death ligand 1 (PD-L1)/programmed cell death ligand 2 (PD-L2)/Janus kinase 2 (JAK2) overexpression; MSI tumors have high mutation and DNA methylation rates; CIN tumors contain chromosomal alterations affecting mainly tyrosine kinase receptors; and finally, GS tumors are chromosomally stable and have a high frequency of cadherin 1 (*CDH1*) and Ras homolog family member A (*RHOA*) mutations [Table 1]^[4]. This review will focus on EBV-positive tumors.

EBV is a double-stranded DNA virus that belongs to the herpesvirus family. An estimated 90% of the human population show signs of previous infection with EBV^[7]. The virus is usually transmitted orally through saliva. Primary infection is most commonly asymptomatic, but it can lead to acute mononucleosis in a subset of the population, primarily adolescents and young adults^[8]. Following primary infection, the virus establishes a lifelong latent infection in the host. EBV can remain latent in both lymphocytes and epithelial cells, where it expresses only a subset of its genes. Depending on which viral genes are expressed, latent EBV infection is typically classified into four latency programs, known as latency 0, I, II, and III [Table 2]^[7].

In 1964, EBV was the first human virus to be associated with cancer, when it was discovered in Burkitt's Lymphoma^[9,10]. Since then, EBV has also been linked to other lymphomas, including Hodgkin lymphoma, diffuse large B-cell lymphoma (DLBCL), and Natural Killer/T-cell lymphomas (NK-T lymphomas)^[11]. In addition, EBV has been associated with certain epithelial cancers, notably Nasopharyngeal Carcinoma (NPC) and GC. In both lymphoid and epithelial cancers, EBV persists in a latent state in infected cells^[11]. However, different EBV-associated cancers demonstrate different viral gene expression patterns [Table 2].

Table 2. Viral genes expressed in each latency program

Latency program	EBV gene expression pattern	Cancer*
0	No viral proteins expressed	
I	EBER 1 & 2, BART miRNAs, EBNA1 <i>Latency I</i> +/- LMP2A	BL GC
II	<i>Latency I</i> + LMP1, LMP2A, LMP2B	NPC, NK-T lymphomas, HL
III	<i>Latency II</i> + EBNA2, EBNA3s, EBNA-LP, BHRF1 miRNAs	PTDL, AIDS-related lymphomas (IB-DLBCL)

*This column indicates some cancers that are typically associated with each latency program. Within each cancer type (e.g., BL and DLBCL) there might be subtypes that show different patterns of latent expression. EBER: Epstein-Barr encoding region; BART: BamHI A rightward transcripts; miRNA: microRNA; BHRF1: BamHI fragment H rightward open reading frame 1; EBNA1: EBV nuclear antigen; LMP: latent membrane protein; EBNA-LP: EBNA leader protein; BL: Burkitt's lymphoma; GC: gastric cancer; NPC: nasopharyngeal carcinoma; DLBCL: diffuse large B-cell lymphoma; NK-T lymphomas: natural killer/T-cell lymphomas; HL: Hodgkin lymphoma; PTDL: post-transplant lymphoproliferative disorder; IB-DLBCL: immunoblastic DLBCL; EBV: Epstein-Barr virus

In EBV-associated GC (EBVaGC), the virus expresses all latency I products, meaning EBV nuclear antigen 1 (EBNA1), Epstein-Barr encoding region small noncoding RNAs, BamHI A rightward transcripts (BARTs), and BART miRNAs, while around 50% of the cases also show Latent Membrane Protein 2A (LMP2A) expression, which is typically associated with latency II^[12]. LMP1, another latency II protein, is typically not detected in EBVaGC samples^[12-14]. There is strong geographical variation in the prevalence of EBVaGC but overall around 10% of gastric adenocarcinomas worldwide are classified as EBV-positive^[15].

In EBVaGC, the EBV genome is mainly maintained as a nuclear episome and cancer cells within the tumor show ubiquitous and monoclonal EBV infection^[12,16]. The monoclonality of EBV infection suggests the clonal selection of virus-infected cells in early stages of cancer development. The role of EBV in gastric carcinogenesis is still under investigation, but EBV infection is thought to contribute to GC progression or maintenance, both directly through the activity of viral proteins or RNAs and indirectly through the induction of somatic alterations in the host genome and epigenome^[17].

PD-L1 EXPRESSION IN EBVAGC

Multiple studies have shown that EBVaGC is commonly characterized by high lymphocytic infiltration in the tumor microenvironment, coupled with overexpression of immune-related genes, including *PD-L1* (also known as *CD274*)^[4,5,18]. *PD-L1* is a glycoprotein, expressed by both cancer cells and stromal immune cells in the tumor, that engages the programmed cell death 1 (PD-1) receptor expressed on the surface of infiltrating cytotoxic T cells (CTLs)^[19]. The interaction between *PD-L1* and PD-1 leads to the inhibition of the tumor-infiltrating CTLs, preventing them from attacking and eliminating tumor cells. *PD-L1* is only one of multiple immune checkpoint genes that are known to be upregulated in EBV-positive compared to EBV-negative cancers. *PD-L2*, Lymphocyte activation gene-3 (*LAG3*), T cell immunoglobulin and mucin domain (*Tim-3*), Cluster of differentiation 80 (*CD80*), Cluster of differentiation 86 (*CD86*), and Indoleamine 2, 3-dioxygenase 1 (*IDO1*) are also upregulated, but *PD-L1* has received particular interest because the PD-1/PD-L1 axis is the target of some recent breakthrough cancer therapies^[20]. Monoclonal antibodies that block the interaction between *PD-L1* and PD-1, thus restoring the ability of the immune system to surveil and attack the tumor, have shown promising results as therapeutic agents against multiple cancers, including non-small cell lung cancer and melanoma^[19]. Recently, the Food and Drug Administration (FDA) approved pembrolizumab, a mAb targeting PD-1, as a third-line therapy for advanced gastric tumors that are positive for *PD-L1* expression based on immunohistochemical (IHC) staining^[21,22].

The clinical efficacy and adverse effects of PD-1/PD-L1 therapy vary tremendously among patients. High expression of *PD-L1* in the tumor has been implicated as a significant predictive biomarker for positive response to PD-1/PD-L1 therapy^[23]. However, several clinical studies have demonstrated that some tumors with high *PD-L1* expression do not respond to PD-1/PD-L1 therapy, while some tumors with moderate or

low PD-L1 expression do show beneficial responses. This discrepancy could be attributed in part to the fact that direct determination of PD-L1 expression in the tumor, which is typically performed by IHC assays, has been proven to be difficult and inconsistent^[24,25]. Understanding the molecular mechanisms that control PD-L1 expression in cancer may give rise to more accurate biomarkers for positive response to PD-1/PD-L1 therapy. In addition, regulators of PD-L1 expression could serve as targets for potential combination therapies.

In a phase II clinical trial, Kim *et al.*^[6] identified EBV-positivity and high mutational load as reliable and independent biomarkers for the clinical efficacy of pembrolizumab in GC patients and recommended considering pembrolizumab as an early therapeutic option for EBVaGC. MSI-high (MSI-H) tumors are characterized by high rates of somatic mutations, resulting in increased presentation of neoantigens and thus stimulation of anti-tumor immunity. As is the case in EBV-positive GCs, MSI-H tumors have high levels of tumor-infiltrating lymphocytes (TILs)^[26]. The FDA has already approved the use of front-line pembrolizumab monotherapy in advanced MSI-H solid tumors of any origin, including stomach^[27].

EBVaGC appears to employ some common and some unique mechanisms for PD-L1 regulation. This review presents the most recent findings on PD-L1 regulation in EBVaGC and discusses some of the discrepancies in the literature, parallels with other EBV-associated cancers, and questions to be addressed in the future [Figure 1].

EVOLUTIONARY PRESSURE FOR PD-L1 OVEREXPRESSION

EBVaGC shows high levels of TILs and thus is under strong evolutionary pressure for the development of immune evasion strategies^[4]. One such strategy is PD-L1 overexpression, which is evident in both cancer and stromal immune cells. It is speculated that the high immune activity in the EBVaGC microenvironment reflects the strong immunogenicity of EBV in the body. Up to 5% of the circulating CD8⁺ T cells in EBV-infected individuals are believed to be reactive to lytic or latent EBV antigens^[28]. Indeed, several groups have detected expression of a subset of lytic genes in EBVaGC^[29-31]. According to Borozan *et al.*^[29], the lytic expression pattern detected in EBVaGC, which includes subsets of both early and late lytic proteins, does not indicate lytic, or abortive lytic, replication. However, the presentation of lytic viral antigens by infected cancer cells might be driving further immune activation. Camargo *et al.*^[32] detected higher levels of circulating antibodies targeting both latent and lytic proteins in patients with EBV-positive compared to EBV-negative GCs, even though virtually all patients in the study were seropositive for antibodies against EBNA and the viral capsid. This further suggests that lytic proteins are expressed in EBVaGC and potentially contribute to the activation of the host immune response.

SOMATIC STRUCTURAL VARIATIONS

Somatic genomic alterations include short variations, such as single nucleotide substitutions and short insertions/deletions (indels), as well as long variations, also known as structural variations (SVs)^[33]. SVs affect large chromosomal regions (longer than 50 base-pairs) and include amplifications, deletions, inversions, and translocations^[33]. Somatic changes occur continuously in the life of an individual and are usually repaired, but their accumulation over time can contribute to carcinogenesis^[33]. EBVaGC only appears in a small percentage of EBV-infected individuals and typically long after primary EBV infection suggesting that somatic genomic changes are likely required for cancer development^[34].

Gene amplification

Recent high-throughput sequencing studies have revealed different genomic alterations that lead to PD-L1 overexpression in EBVaGC. TCGA performed a somatic copy-number aberrations analysis in gastric adenocarcinoma samples and identified a frequent (15% of EBVaGC cases) somatic focal amplification at the chromosomal region 9p24.1, which includes the genes *PD-L1*, *PD-L2*, and *JAK2*^[4]. This amplification




Figure 1. The mechanisms of PD-L1 regulation in EBVaGC discussed in this review. EBV: Epstein-Barr virus; EBVaGC: EBV-associated gastric cancer; PD-L1: programmed cell death ligand 1; IFN γ : interferon gamma; TNF α : tumor necrosis factor alpha; EBER: Epstein-Barr encoding region; JAK: Janus kinase; STAT: signal transducers and activators of transcription; IRF: interferon regulatory factor 1; EBNA1: EBV nuclear antigen 1; BART: BamHI A rightward transcripts; PIAS: protein inhibitor of activated STAT 3; I κ B α : nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha; NF κ B: nuclear factor kappa-light-chain-enhancer of activated B cells; LMP2A: latent membrane protein 2A; PTEN: phosphatase and tensin homolog; PIK3CA: phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha; PI3K: phosphoinositide 3-kinase; AKT: protein kinase B; ARID1A: AT-rich interaction domain 1A; mTOR: mammalian target of rapamycin; 3'-UTR: 3' untranslated region; SV: structural variation; SNV: single nucleotide variation; BHRF1: BamHI fragment H rightward open reading frame 1. Created with BioRender.com

was significantly more common in EBV-positive than EBV-negative tumors. Importantly, they showed that the 9p24.1 amplification correlated positively with *PD-L1* mRNA levels in the tumor^[4]. Saito *et al.*^[35] detected 9p24.1 amplification events at a similar rate in a different EBVaGC cohort and demonstrated a direct association between *PD-L1* copy number and protein expression. Using fluorescence in-situ hybridization and IHC staining, they showed that *PD-L1* genomic amplifications were specifically detected in cancer cells that showed high PD-L1 protein expression. Similar amplification patterns have been detected in other cancers, including hepatocellular carcinomas (HCC) and DLBCL^[23,36].

Integration of EBV to the host genome

In EBVaGC, the EBV genome is mostly maintained in the nucleus as a circular episome^[16]. Some studies have also reported viral integration events, even though they are considered rare^[4,37,38]. TCGA identified the presence of RNA sequencing (RNAseq) reads corresponding to a fusion between the host *PLGRKT* gene and the EBV gene *BHLF1* in one EBVaGC sample (FP-7998-01)^[4]. This is notable given that the *PLGRKT* lies within the 9p24.1 chromosomal region and is adjacent to the *PD-L1* locus. It was later shown in the same sample, that the amplified region containing the integrated EBV genome (Copy Number = 4) has a breakpoint in the *PD-L1* 3' untranslated region (3'-UTR)^[37]. Xu *et al.*^[38] performed a genome-wide EBV integration analysis in multiple malignancies and identified that EBV integrates primarily in unstable

chromosomal regions of the host genome. They identified EBV integration events in 25% (10/39) of the GC samples analyzed, with some of the integration breakpoints mapping close to known tumor suppressor genes^[38]. None of their samples had integrated virus close to *PD-L1*, but studies with bigger EBVaGC sample sizes are necessary to identify the frequency and significance of viral integration in or close to the *PD-L1* locus. Viral integration in the host genome has been associated with increased PD-L1 expression in other virus-associated cancers. Cao *et al.*^[20] identified integrated HPV genomes in the *PD-L1* or *PD-L2* loci in three cases from the TCGA Head and Neck Squamous Cell Carcinoma cohort and showed that these integration events correlated with elevated PD-L1 and PD-L2 expression.

3'-UTR structural variations

SVs in the *PD-L1* 3'-UTR have also been associated with increased PD-L1 expression in EBVaGC^[34,37]. Kataoka *et al.*^[37] analyzed RNAseq data from all TCGA cancer types and searched for 3'-UTR disruptions in *PD-L1*. The authors identified *PD-L1* 3'-UTR truncations in 31/10,210 cancer cases and showed that they correlated with high PD-L1 expression. The highest frequency of 3'-UTR truncations was found in DLBCL (4/48) and GC (9/415), with a third of the GC samples (3/9) being EBV-positive. Therefore, around 10% of EBVaGC samples in TCGA were found to have *PD-L1* 3'-UTR SVs^[34,37]. In a follow-up study, Kataoka *et al.*^[34] analyzed samples from multiple EBV-associated lymphomas and found that *PD-L1* 3'-UTR SVs were significantly more common in EBV-positive compared to EBV-negative lymphomas. They report that *PD-L1* 3'-UTR genomic truncations in cell lines and mouse models promote PD-L1 overexpression and immune evasion, consistent with the patient data^[37].

POST-TRANSCRIPTIONAL REGULATION

The 3'-UTR contains sequences or structural regions, called regulatory elements, that are important for the post-transcriptional regulation of a gene. These regulatory elements control binding to miRNAs and RNA-binding proteins (RBP), which influence mRNA stability, translation rate, and localization^[39]. miRNAs are short non-coding RNAs that silence gene expression by binding to complementary sequences in the 3'-UTR of target mRNAs. miRNA-mRNA binding usually triggers mRNA degradation or blocks translation. The fact that 3'-UTR shortening has such a profound effect on PD-L1 expression in multiple cancers indicates that PD-L1 is under tight post-transcriptional control^[37].

3'-UTR short variations

Mutations in the 3'-UTR have the capacity to remove existing or create new binding sites for miRNAs and RBPs. Some germline and somatic mutations in the 3'-UTR of *PD-L1* have been shown to correlate with PD-L1 expression in gastric and other cancers^[40-44]. Wu *et al.*^[43] analyzed 728 GC samples and found that the AA and AG genotypes in rs2297136, a germline single nucleotide polymorphism (SNP) located in the 3'-UTR of *PD-L1*, were associated with lower PD-L1 protein levels. They reported that the miRNAs miR-324-5p and miR-362 are predicted to bind to that region of the *PD-L1* 3'-UTR, but no validation experiments were pursued. Wang *et al.*^[44] polymerase chain reaction (PCR)-amplified and sequenced the 3'-UTR of *PD-L1* in hundreds of GC and matched normal samples and identified a frequent guanine-to-cytosine somatic mutation that correlated with increased PD-L1 protein expression. It was shown that this mutation maps to a seed-binding region for miR-570 and it was proven experimentally that it increases PD-L1 expression by disrupting miR-570 binding^[44]. To date, most studies looking at *PD-L1* 3'-UTR mutations have been low-throughput, with small sample sizes or targeted on specific SNP locations. There has not been a comprehensive study looking at the frequency and effect of all possible somatic and germline variants in the *PD-L1* 3'-UTR in EBVaGC or other EBV-associated cancers. The fact that SVs in the *PD-L1* 3'-UTR appear to occur more frequently in EBV-positive than EBV-negative cancers raises the question of whether short variants in the 3'-UTR could be an alternative or parallel mechanism for PD-L1 overexpression. Large-scale variant calling studies in gastric and other cancers, including the TCGA somatic mutation data, have mostly relied on whole exome sequencing data and exclude 3'-UTR sequences. This has created a gap in our understanding of 3'-UTR variations in cancer in general.

Host miRNAs and RBPs

Multiple miRNAs have been implicated in the control of PD-L1 expression in GC^[45]. Some miRNAs, such as miR-152 and miR-200, target the 3'-UTR of PD-L1 directly. Other miRNAs, such as miR-19a and miR-19b, affect PD-L1 levels indirectly, by targeting signaling pathways or transcription factors that regulate PD-L1 expression^[45-48].

EBVaGC has been reported to exhibit a distinct host miRNA expression profile from other GC subtypes^[4,49,50]. Differentially expressed miRNAs include ones that have been independently shown to target PD-L1, with the most notable example being miR-200^[48]. miR-200 is a family of miRNAs found in two distinct genomic clusters and consists of miR-200a, miR-200b, miR-429, miR-200c, and miR-141^[51]. miR-200a, miR-200b, and miR-429 form a cluster on chromosome 1, while miR-200c and miR-141 form a cluster on chromosome 12. miR-200b, miR-200c and miR-429 share the same seed sequence, while miR-200a and miR-141 have a seed sequence that differs from the others by one nucleotide. The *PD-L1* 3'-UTR contains one binding site for each seed-sharing functional cluster of miR-200 miRNAs and all miR-200a, b, and c have been shown to directly silence PD-L1 expression^[48].

The miR-200 family has been shown to be downregulated in EBV-positive compared to EBV-negative GC samples, as well as in EBV-negative GC cells following infection with recombinant EBV (rEBV) *in vitro*^[52,53]. Whether or how EBV latent proteins downregulate miR-200 remains unclear. Shinozaki *et al.*^[53] reported that overexpression of any of *EBNA1*, *LMP2A*, or *BARF0* (a *BART* transcript) in EBV-negative GC cells (MKN74 cell line) leads to miR-200a/b transcriptional repression, while overexpression of EBERs downregulates the mature miRNAs post-transcriptionally. The authors concluded that EBV latent proteins and RNAs act synergistically to downregulate miR-200. This is in contrast with a study by Marquitz *et al.*^[52] on a different EBV-negative GC cell line, AGS. They reported that rEBV-infected AGS (AGS-EBV) cells showed consistent downregulation in some tumor suppressor miRNAs, including miR-200, when compared to the parental uninfected cells. However, *EBNA1* or *LMP1* overexpression in AGS did not affect the expression levels of miR-200 and most of the other EBV-downregulated miRNAs^[52]. They speculate that EBV-mediated cellular miRNA downregulation might not be mediated by the latent viral proteins, but by viral miRNAs, through their effect on host transcription factors, or by EBERs^[52]. Another possible mechanism of host miRNA downregulation in EBV-infected cells is methylation of their promoter sequences. EBV infection in GC cells has been shown to promote extensive DNA methylation, partly through the activity of the latent protein LMP2A, which induces DNA methyltransferases (DNMTs)^[54,55]. Indeed, the miR-200c promoter contains CpG islands that have been shown to be methylated by DNMT3a in GC^[56].

Research from our group described an EBV gene/host miRNA/PD-L1 regulation axis in B cell lymphomas, where the EBV protein EBNA2 induces PD-L1 expression by transcriptionally repressing miR-34a, a miRNA that downregulates PD-L1^[57]. EBNA2 is a latency III-associated protein and thus is not expressed in GC. However, miR-34a has been shown to be downregulated in GC overall and to be transcriptionally repressed by EBNA1 in EBVaGC^[58,59].

The *PD-L1* 3'-UTR contains multiple adenosine-uridine (AU)-rich elements (AREs) that are known to serve as binding sites for different RBPs^[60]. In GC, the RBP tristetraprolin (TTP) has been shown to bind to AREs in the *PD-L1* 3'-UTR and to promote mRNA destabilization, leading to reduced PD-L1 expression^[62]. In addition, in NPC, the EBV latent protein LMP1 reduces TTP expression through extracellular-signal-regulated kinase 1/2 (ERK1/2) activation^[63]. Nonetheless, there has not been an association between EBV infection and TTP expression in GC.

Viral miRNAs

EBV expresses 25 precursor miRNAs (pre-miRNAs) and 44 mature miRNAs. Three pre-miRNAs are derived from the BamHI fragment H rightward open reading frame 1 (BHRF1) region of the viral genome,

while the rest are derived from the BART region^[64]. BART miRNAs are thought to play an important role in EBVaGC and are highly abundant, accounting for up to 15-20% of the total miRNA pool in some EBVaGC cell lines^[65]. Some studies have demonstrated that xenografts of EBV-infected GC cell lines, including AGS-EBV and SNU719 cells, in immunocompromised mice show up to 10- or 100-fold overexpression of BART miRNAs compared to their parental cell lines^[66,67]. This suggests that BART miRNAs might be particularly important for cancer progression *in vivo*. BART miRNAs are known to play a range of roles in GC, targeting host cell apoptosis, cell cycle, and metastasis^[68,69]. Their role in immune evasion is less well studied in GC, although multiple immune-related functions have been attributed to BART miRNAs in other EBV-associated cancers^[64].

In contrast to BART miRNAs, which are expressed in all EBV latent stages, BHRF1 miRNAs are thought to be expressed only in latency III and lytic replication. BHRF1 miRNAs are commonly reported as barely detectable in GC tissues and cell lines^[4,54,62]. However, Treece *et al.*^[49] analyzed miRNA expression in FFPE tissues from 78 cases of gastric adenocarcinoma, including 20 EBVaGCs, and reported that BHRF1-2-5p was significantly overexpressed in EBV-infected vs. EBV-negative cancers, although to a lower extent than BART miRNAs. Marquitz *et al.*^[52] also detected low expression of BHRF1 miRNAs in sequencing from EBV-infected GC cells (AGS-EBV). Both groups attributed the detection of BHRF1 miRNAs to possible low levels of viral replication. Given that several studies have reported the expression of a subset of lytic transcripts in EBVaGC, even in the absence of lytic replication, the level and importance of BHRF1 miRNA expression remain to be determined. In an *in vitro* model of EBV-driven B-cell differentiation, BHRF1-2-5p was found to downregulate PD-L1^[70]. The same study identified potential binding sites for some BART miRNAs in the *PD-L1* 3'-UTR, including BART19-3p, but overexpression of the miRNA did not appear to influence PD-L1 expression. Four BART miRNAs, BART2-5p, BART7-3p, BART14-3p, miR-BART22 were found to interact with *PD-L1* mRNA in a high-throughput photoactivatable ribonucleoside-enhanced crosslinking and immunoprecipitation (PAR-CLIP) study in lymphoblastoid cell lines, but no validation experiments have been performed^[71,72]. A recent study reported that miR-BART5-5p, which shares seed homology with the host miRNAs miR-18a-5p and miR-18b-5p, leads to signal transducers and activators of transcription 3 (STAT3)-dependent transcriptional PD-L1 upregulation by targeting the STAT3 inhibitor Protein Inhibitor Of Activated STAT 3 (PIAS3)^[73].

INTRINSIC SIGNALING

PD-L1 expression can be dysregulated by oncogenic activation of signaling pathways like the JAK/STAT, phosphoinositide 3-kinase (PI3K)/protein kinase B (Akt)/mammalian target of rapamycin (mTOR), Mitogen-activated protein kinase/ERK kinase (MEK)/ERK, and Jun/Activator protein 1 (AP-1) pathways. These pathways act independently or synergistically to control PD-L1 expression, at the transcriptional, post-transcriptional, and post-translational stage^[74-76]. Their importance in PD-L1 regulation tends to vary among different cancer types. The constitutive activation of an intrinsic signaling pathway in cancer is usually the result of mutations or SVs in genes of key components or regulators of the pathway. In the case of virus-associated cancers, viral proteins can also induce constitutive signaling in infected cells.

Signaling activation by host gene mutations

A frequently mutated gene in EBVaGC cancer is *PIK3CA*, which encodes a catalytic component of the PI3K kinase^[4,77]. The most common variants in *PIK3CA* are associated with increased PI3K signaling activity^[77]. The PI3K pathway is thought to regulate PD-L1 expression in a tissue-specific manner. Loss of Phosphatase and tensin homolog (PTEN) leads to activation of the PI3K pathway and induction of PD-L1 expression in gliomas and colorectal cancer^[78,79]. In gliomas, Parsa *et al.*^[79] showed that the PI3K/Akt/mTOR pathway increases *PD-L1* mRNA translation through polysomal recruitment. In the case of GC, there is conflicting evidence for the importance of the PI3K pathway in PD-L1 regulation. Supporting the importance of PI3K signaling in promoting PD-L1 expression, Kim *et al.*^[80] showed that the PI3K inhibitor

LY294002 reduced PD-L1 expression in three GC cell lines. In addition, Menyhárt *et al.*^[81] performed hierarchical clustering to determine the mutations that could best stratify TCGA GC patients based on PD-L1 expression. They showed that the mutation status of *PIK3CA* served as the best root node for the stratification, while mutations in other immune-related genes like *MEF2C*, *SLC11A1*, and *KIF15* could help further refine it, suggesting potential interactions between these genes for the control of PD-L1 expression^[81]. However, Mimura *et al.*^[82] performed experiments with a panel of GC cell lines and determined that the PI3K inhibitor wortmannin did not affect PD-L1 expression. Besides, Seo *et al.*^[77] analyzed 112 EBV-positive GC samples and reported that *PIK3CA* mutations did not show any correlation with PD-L1 expression or TIL abundance. Therefore, the importance of *PIK3CA* mutations and PI3K signaling in the control of PD-L1 expression in GC and EBVaGC is still unclear, and may be dependent on interactions with other mutations and signaling pathways.

Another gene that is commonly mutated in EBVaGC is AT-Rich Interaction Domain 1A (*ARID1A*), which encodes an important component of the SWItch/Sucrose Non-Fermentable (SWI/SNF) chromatin remodeling complex^[4]. Mutations in *ARID1A* are usually loss-of-function mutations. Kim *et al.*^[80] analyzed 273 GC samples and showed that *ARID1A* protein loss correlated with PD-L1 positivity (as defined by IHC staining), independently of EBV or MSI status. They also showed that *ARID1A* knockdown *in vitro* directly leads to PD-L1 overexpression, through the activation of the PI3K/Akt pathway. To account for the variability in PD-L1 expression among *ARID1A*-mutated tumors, the authors looked for additional mutations that could be acting synergistically with *ARID1A* mutations. They found *KRAS* mutations in the three *ARID1A*-mutated MSI-H tumors with the highest PD-L1 expression, while two other samples that harbored *KRAS* but not *ARID1A* mutations did not show elevated PD-L1 levels. These data further suggest that PD-L1 expression is controlled by multiple oncogenic signaling pathways acting in coordination.

Signaling activation by viral latent proteins

The EBV latent membrane proteins (LMPs) are potent oncoproteins and are known to activate oncogenic signaling cascades in EBV-associated malignancies. In epithelial cancers, LMP2A has been found to regulate the nuclear factor kappa-light-chain-enhancer of activated B cells (NFκB), PI3K/Akt, MEK/ERK, and transforming growth factor beta (TGFβ) pathways^[83-86]. In GC specifically, LMP2A has been reported to lead to constitutive NFκB activation by inhibiting the expression of the nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (IκBα), which negatively regulates NFκB activity^[83,84]. The mechanism of IκBα downregulation appears to be unrelated to promoter methylation and remains to be determined. Independent studies have shown that NFκB promotes PD-L1 transcription in GC^[87-89], but no direct association has been made between LMP2A, NFκB, and PD-L1. As mentioned above, LMP2A is also thought to be important for promoting the DNA hypermethylated state observed in EBVaGC. Hino *et al.*^[90] showed that LMP2A induces phosphorylation and activation of STAT3, which in turn leads to overexpression of DNMT1, thus changing the DNA methylation landscape of the cell. They showed that one of the targets of DNMT1 is *PTEN*, which encodes a negative regulator of the PI3K pathway. The hypermethylation of the *PTEN* promoter reduces *PTEN* expression and leads to overactivation of the PI3K pathway^[90]. Interestingly, *PTEN* is also known to be directly targeted by miR-BART1 and BART7-3p in NPC^[91,92]. Moon *et al.*^[93] reported that the small interfering RNA (siRNA)-mediated knockdown of LMP2A did not affect PD-L1 expression in SNU719, an EBVaGC cell line. However, more studies with additional methods of manipulating LMP2A expression and activity are required to elucidate its role in PD-L1 regulation.

INTERFERON GAMMA-INDUCIBLE PD-L1 EXPRESSION

Multiple studies have reported that GC patients with higher levels of CD8⁺ TILs also have higher PD-L1 expression^[94-96]. This suggests that PD-L1 overexpression in EBVaGC is at least partly a result of the evolutionary pressure from the adaptive immune response acting against the cancer. When CD8⁺ T cells are

activated, such as through the recognition of neoantigens or viral antigens presented by cancer cells, they produce inflammatory cytokines like tumor necrosis factor alpha (TNF α) and interferon gamma (IFN γ)^[97,98]. IFN γ has been shown to induce PD-L1 expression in various cancers^[99]. In GC, PD-L1 induction by IFN γ appears to occur mainly through the JAK/STAT/Interferon regulatory factor 1 (IRF1) signaling axis^[82,100-102]. The extent to which different cancer cells are responsive to IFN γ and the downstream effects of IFN γ exposure vary among cancer types and molecular subgroups within a cancer type.

In TCGA and other GC cohorts, when EBVaGC samples are compared to EBV-negative ones, they demonstrate elevated IFN γ signatures, indicated by higher expression of IFN γ , JAK/STAT signaling components, and several IFN γ -induced genes^[4,103]. In addition, *in vitro* studies have shown that EBVaGC cell lines induce PD-L1 expression in response to IFN γ to a much higher extent than EBV-negative GC cell lines^[93,104]. PD-L1 induction in response to IFN γ has also been shown to be significantly elevated in other EBV-associated epithelial malignancies, such as NPC^[105]. In NPC, the viral protein LMP1 acts synergistically with IFN γ to induce PD-L1 expression through the activation of the JAK3/STAT3, NF κ B, and AP-1 signaling pathways. In GC, EBNA1 has been shown to promote IFN γ -induced PD-L1 overexpression. Moon *et al.*^[93] showed that *EBNA1* knockdown in SNU719, an EBVaGC cell line, resulted in the transcriptional downregulation of JAK2. *EBNA1* knockdown also resulted in a small but significant reduction in constitutive and IFN γ -induced PD-L1 levels. However, ectopic EBNA1 expression in the EBV-negative GC cell line AGS did not affect constitutive or IFN γ -induced PD-L1 expression. In contrast, Su *et al.*^[106] reported that AGS-EBV cells are more sensitive to IFN γ /TNF α treatment, showing higher downstream PD-L1 upregulation, than the uninfected parental cell line. This suggests that other viral factors, in addition to EBNA1, might be necessary for increasing IFN γ -induced PD-L1 expression in EBV-infected GC cells. Further studies are required to determine how EBNA1 and other viral or host factors promote increased IFN γ sensitivity and PD-L1 expression in EBVaGC.

Nakayama *et al.*^[107] analyzed 43 EBVaGC samples and showed that the number of EBV genomes per cancer cell (EBV copy number) correlates positively with PD-L1 expression. Similarly, Strong *et al.*^[31] performed cellular gene expression analysis on 12 EBV-positive and 20 EBV-negative TCGA GC samples and showed that, following hierarchical clustering, the 4 EBV-positive samples that had a much higher EBV coverage depth (in RNAseq data) than the rest of the EBV-positive samples formed a well-defined gene expression cluster. When they compared expression between “high” and “low” EBV GC samples, a large proportion of the genes that were upregulated in the “high” EBVaGC group were immune-related, including IFN γ , STAT1, IRF1, and multiple IFN γ -induced genes^[31]. As the authors state, EBERs, which have been shown to induce IFN γ and TNF α production in peripheral blood mononuclear cells *in vitro*, could be contributing to the elevated IFN γ signature found in “high” compared to “low” EBV GC samples^[108].

CONCLUSION

Even though GC is declining in the United States, it still has one of the lowest 5-year survival rates of any cancer type^[2]. This highlights the need for new therapeutic strategies, especially for metastatic cases that have the poorest prognosis and account for most of the new diagnoses every year. The molecular heterogeneity of GC correlates with the response rate to different therapies, indicating that different approaches should be considered for different molecular subgroups. The FDA recently approved pembrolizumab as second-line therapy for patients with advanced MSI-H tumors of any type, including GC^[109]. In the last few years, several molecular and clinical studies present EBVaGC as another subgroup of GC that could benefit from early-line treatment with immune checkpoint inhibitors^[6,107].

In EBVaGC, high immune activation in the tumor microenvironment likely acts as a driving force for the selection of immune escape mechanisms such as PD-L1 overexpression. Different mechanisms act independently or synergistically to induce PD-L1 expression. These include somatic genomic modifications,

oncogenic activation of intrinsic signaling pathways, increased sensitivity to PD-L1 inducing signals from the tumor microenvironment, and post-transcriptional control mechanisms. Overall, the regulation of PD-L1 expression in EBVaGC is poorly understood and further studies are necessary to explain how EBV and host factors contribute to it. Insight into the important genetic and epigenetic factors that control PD-L1 expression in EBVaGC and other cancers could reveal new biomarkers for positive response to immunotherapy, as well as novel therapeutic targets.

DECLARATIONS

Acknowledgments

We especially appreciate Dr. Maria Mavrikaki, Allison Baker, Tanvi Saxena, and Jason Nomburg for their comments on this manuscript.

Authors' contributions

Wrote the manuscript: Miliotis CN, Slack FJ

Availability of data and materials

Not applicable.

Financial support and sponsorship

Slack FJ acknowledges the NCI Outstanding Investigator Award (R35CA232105) and the support of the Ludwig Institute at Harvard and the V Foundation.

Conflicts of interest

Slack FJ discloses financial interests and SAB roles with Mira DX and MiRNA Therapeutics. The other author declares no competing financial interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Rawla P, Barsouk A. Epidemiology of gastric cancer: global trends, risk factors and prevention. *Prz Gastroenterol* 2019;14:26-38.
2. Street W. Cancer facts & figures 2019. 1930:76.
3. Zali H, Rezaei-Tavirani M, Azodi M. Gastric cancer: prevention, risk factors and treatment. *Gastroenterol Hepatol Bed Bench* 2011;4:175-85.
4. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* 2014;513:202-9.
5. Angell HK, Lee J, Kim KM, Kim K, Kim ST, et al. PD-L1 and immune infiltrates are differentially expressed in distinct subgroups of gastric cancer. *Oncoimmunology* 2019;8:e1544442.
6. Kim ST, Cristescu R, Bass AJ, Kim KM, Odegaard JI, et al. Comprehensive molecular characterization of clinical responses to PD-1 inhibition in metastatic gastric cancer. *Nat Med* 2018;24:1449-58.
7. Rickinson A. Epstein-Barr virus. *Virus Res* 2001;82:109-13.
8. Balfour HH Jr, Dunmire SK, Hogquist KA. Infectious mononucleosis. *Clin Transl Immunol* 2015;4:e33.
9. Epstein M, Achong B, Barr Y. Virus particles in cultured lymphoblasts from Burkitt's lymphoma. *Lancet* 1964;283:702-3.
10. Dalldorf G, Linsell CA, Barnhart FE, Martyn R. An epidemiologic approach to the lymphomas of African children and Burkitt's sarcoma of the jaws. *Perspect Biol Med* 1964;7:435-49.
11. Hsu JL, Glaser SL. Epstein-Barr virus-associated malignancies: epidemiologic patterns and etiologic implications. *Crit Rev Oncol*

- Hematol 2000;34:27-53.
12. Imai S, Koizumi S, Sugiura M, Tokunaga M, Uemura Y, et al. Gastric carcinoma: monoclonal epithelial malignant cells expressing Epstein-Barr virus latent infection protein. *Proc Natl Acad Sci U S A* 1994;91:9131-5.
 13. Ribeiro J, Oliveira A, Malta M, Oliveira C, Silva F, et al. Clinical and pathological characterization of Epstein-Barr virus-associated gastric carcinomas in Portugal. *World J Gastroenterol* 2017;23:7292-302.
 14. Oh ST, Seo JS, Moon UY, Kang KH, Shin DJ, et al. A naturally derived gastric cancer cell line shows latency I Epstein-Barr virus infection closely resembling EBV-associated gastric cancer. *Virology* 2004;320:330-6.
 15. Nogueira C, Mota M, Gradiz R, Cipriano MA, Caramelo F, et al. Prevalence and characteristics of Epstein-Barr virus-associated gastric carcinomas in Portugal. *Infect Agent Cancer* 2017;12:41.
 16. Fukayama M, Hayashi Y, Iwasaki Y, Chong J, Ooba T, et al. Epstein-Barr virus-associated gastric carcinoma and Epstein-Barr virus infection of the stomach. *Lab Invest* 1994;71:73-81.
 17. Naseem M, Barzi A, Brezden-Masley C, Puccini A, Berger MD, et al. Outlooks on Epstein-Barr virus associated gastric cancer. *Cancer Treat Rev* 2018;66:15-22.
 18. Derks S, Liao X, Chiaravalli AM, Xu X, Camargo MC, et al. Abundant PD-L1 expression in Epstein-Barr Virus-infected gastric cancers. *Oncotarget* 2016;7:32925-32.
 19. Salmaninejad A, Valilou SF, Shabgah AG, Aslani S, Alimardani M, et al. PD-1/PD-L1 pathway: Basic biology and role in cancer immunotherapy. *J Cell Physiol* 2019;234:16824-37.
 20. Cao S, Wylie KM, Wyczalkowski MA, Karpova A, Ley J, et al. Dynamic host immune response in virus-associated cancers. *Commun Biol* 2019;2:109.
 21. Fuchs CS, Doi T, Jang RW, Muro K, Satoh T, et al. Safety and efficacy of pembrolizumab monotherapy in patients with previously treated advanced gastric and gastroesophageal junction cancer: phase 2 clinical KEYNOTE-059 Trial. *JAMA Oncol* 2018;4:e180013.
 22. Fashoyin-Aje L, Donoghue M, Chen H, He K, Veeraraghavan J, et al. FDA approval summary: pembrolizumab for recurrent locally advanced or metastatic gastric or gastroesophageal junction adenocarcinoma expressing PD-L1. *Oncologist* 2019;24:103-9.
 23. Ma LJ, Feng FL, Dong LQ, Zhang Z, Duan M, et al. Clinical significance of PD-1/PD-Ls gene amplification and overexpression in patients with hepatocellular carcinoma. *Theranostics* 2018;8:5690-702.
 24. Brar G, Shah MA. The role of pembrolizumab in the treatment of PD-L1 expressing gastric and gastroesophageal junction adenocarcinoma. *Therap Adv Gastroenterol* 2019;12:1756284819869767.
 25. Rimm DL, Han G, Taube JM, Yi ES, Bridge JA, et al. A prospective, multi-institutional, pathologist-based assessment of 4 immunohistochemistry assays for PD-L1 expression in non-small cell lung cancer. *JAMA Oncol* 2017;3:1051-8.
 26. Cho J, Chang YH, Heo YJ, Kim S, Kim NK, et al. Four distinct immune microenvironment subtypes in gastric adenocarcinoma with special reference to microsatellite instability. *ESMO Open* 2018;3:e000326.
 27. Marcus L, Lemery SJ, Keegan P, Pazdur R. FDA approval summary: pembrolizumab for the treatment of microsatellite instability-high solid tumors. *Clin Cancer Res* 2019;25:3753-8.
 28. Hislop AD, Annels NE, Gudgeon NH, Leese AM, Rickinson AB. Epitope-specific evolution of human CD8(+) T cell responses from primary to persistent phases of Epstein-Barr virus infection. *J Exp Med* 2002;195:893-905.
 29. Boroza I, Zapata M, Frappier L, Ferretti V. Analysis of Epstein-Barr virus genomes and expression profiles in gastric adenocarcinoma. *J Virol* 2018;92:e01239-17.
 30. Song H, Lim Y, Im H, Bae JM, Kang GH, et al. Interpretation of EBV infection in pan-cancer genome considering viral life cycle: LiEB (Life cycle of Epstein-Barr virus). *Sci Rep* 2019;9:3465.
 31. Strong MJ, Xu G, Coco J, Baribault C, Vinay DS, et al. Differences in gastric carcinoma microenvironment stratify according to EBV infection intensity: implications for possible immune adjuvant therapy. *PLoS Pathog* 2013;9:e1003341.
 32. Camargo MC, Kim KM, Matsuo K, Torres J, Liao LM, et al. Circulating antibodies against Epstein-Barr virus (EBV) and p53 in EBV-positive and -negative gastric cancer. *Cancer Epidemiol Biomarkers Prev* 2020;29:414-9.
 33. Yi K, Ju YS. Patterns and mechanisms of structural variations in human cancer. *Exp Mol Med* 2018;50:98.
 34. Kataoka K, Miyoshi H, Sakata S, Dobashi A, Couronné L, et al. Frequent structural variations involving programmed death ligands in Epstein-Barr virus-associated lymphomas. *Leukemia* 2019;33:1687-99.
 35. Saito R, Abe H, Kunita A, Yamashita H, Seto Y, et al. Overexpression and gene amplification of PD-L1 in cancer cells and PD-L1+ immune cells in Epstein-Barr virus-associated gastric cancer: the prognostic implications. *Mod Pathol* 2017;30:427-39.
 36. Wang Y, Wenzl K, Manske MK, Asmann YW, Sarangi V, et al. Amplification of 9p24.1 in diffuse large B-cell lymphoma identifies a unique subset of cases that resemble primary mediastinal large B-cell lymphoma. *Blood Cancer J* 2019;9:73.
 37. Kataoka K, Shiraishi Y, Takeda Y, Sakata S, Matsumoto M, et al. Aberrant PD-L1 expression through 3'-UTR disruption in multiple cancers. *Nature* 2016;534:402-6.
 38. Xu M, Zhang WL, Zhu Q, Zhang S, Yao YY, et al. Genome-wide profiling of Epstein-Barr virus integration by targeted sequencing in Epstein-Barr virus associated malignancies. *Theranostics* 2019;9:1115-24.
 39. Mayr C. Regulation by 3'-untranslated regions. *Annu Rev Genet* 2017;51:171-94.
 40. Nomizo T, Ozasa H, Tsuji T, Funazo T, Yasuda Y, et al. Clinical impact of single nucleotide polymorphism in PD-L1 on response to nivolumab for advanced non-small-cell lung cancer patients. *Sci Rep* 2017;7:45124.
 41. Yeo MK, Choi SY, Seong IO, Suh KS, Kim JM, et al. Association of PD-L1 expression and PD-L1 gene polymorphism with poor prognosis in lung adenocarcinoma and squamous cell carcinoma. *Hum Pathol* 2017;68:103-11.
 42. Lee SY, Jung DK, Choi JE, Jin CC, Hong MJ, et al. Functional polymorphisms in PD-L1 gene are associated with the prognosis of

- patients with early stage non-small cell lung cancer. *Gene* 2017;599:28-35.
43. Wu Y, Zhao T, Jia Z, Cao D, Cao X, et al. Polymorphism of the programmed death-ligand 1 gene is associated with its protein expression and prognosis in gastric cancer. *J Gastroenterol Hepatol* 2019;34:1201-7.
 44. Wang W, Sun J, Li F, Li R, Gu Y, et al. A frequent somatic mutation in CD274 3'-UTR leads to protein over-expression in gastric cancer by disrupting miR-570 binding. *Hum Mutat* 2012;33:480-4.
 45. Wang Q, Lin W, Tang X, Li S, Guo L, et al. The roles of microRNAs in regulating the expression of PD-1/PD-L1 immune checkpoint. *Int J Mol Sci* 2017;18:2540.
 46. Wang F, Li T, Zhang B, Li H, Wu Q, et al. MicroRNA-19a/b regulates multidrug resistance in human gastric cancer cells by targeting PTEN. *Biochem Biophys Res Commun* 2013;434:688-94.
 47. Wang Y, Wang D, Xie G, Yin Y, Zhao E, et al. MicroRNA-152 regulates immune response via targeting B7-H1 in gastric carcinoma. *Oncotarget* 2017;8:28125-34.
 48. Chen L, Gibbons DL, Goswami S, Cortez MA, Ahn YH, et al. Metastasis is regulated via microRNA-200/ZEB1 axis control of tumour cell PD-L1 expression and intratumoral immunosuppression. *Nat Commun* 2014;5:5241.
 49. Treece AL, Duncan DL, Tang W, Elmore S, Morgan DR, et al. Gastric adenocarcinoma microRNA profiles in fixed tissue and in plasma reveal cancer-associated and Epstein-Barr virus-related expression patterns. *Lab Invest* 2016;96:661-71.
 50. Alessandrini L, Manchi M, De Re V, Dolcetti R, Canzonieri V. Proposed molecular and miRNA classification of gastric cancer. *Int J Mol Sci* 2018;19:1683.
 51. Humphries B, Yang C. The microRNA-200 family: small molecules with novel roles in cancer development, progression and therapy. *Oncotarget* 2015;6:6472-98.
 52. Marquitz AR, Mathur A, Chugh PE, Dittmer DP, Raab-Traub N. Expression profile of microRNAs in Epstein-Barr virus-infected AGS gastric carcinoma cells. *J Virol* 2014;88:1389-93.
 53. Shinozaki A, Sakatani T, Ushiku T, Hino R, Isogai M, et al. Downregulation of microRNA-200 in EBV-associated gastric carcinoma. *Cancer Res* 2010;70:4719-27.
 54. Zhao J, Liang Q, Cheung KF, Kang W, Lung RW, et al. Genome-wide identification of Epstein-Barr virus-driven promoter methylation profiles of human genes in gastric cancer cells. *Cancer* 2013;119:304-12.
 55. Queen KJ, Shi M, Zhang F, Cvek U, Scott RS. Epstein-Barr virus-induced epigenetic alterations following transient infection. *Int J Cancer* 2013;132:2076-86.
 56. Li Y, Nie Y, Tu S, Wang H, Zhou Y, et al. Epigenetically deregulated miR-200c is involved in a negative feedback loop with DNMT3a in gastric cancer cells. *Oncol Rep* 2016;36:2108-16.
 57. Anastasiadou E, Stroopinsky D, Alimperti S, Jiao AL, Pyzer AR, et al. Epstein-Barr virus-encoded EBNA2 alters immune checkpoint PD-L1 expression by downregulating miR-34a in B-cell lymphomas. *Leukemia* 2019;33:132-47.
 58. Zhou Y, Ding BZ, Lin YP, Wang HB. MiR-34a, as a suppressor, enhance the susceptibility of gastric cancer cell to luteolin by directly targeting HK1. *Gene* 2018;644:56-65.
 59. Kim SM, Hur DY, Hong SW, Kim JH. EBV-encoded EBNA1 regulates cell viability by modulating miR34a-NOX2-ROS signaling in gastric cancer cells. *Biochem Biophys Res Commun* 2017;494:550-5.
 60. Otsuka H, Fukao A, Funakami Y, Duncan KE, Fujiwara T. Emerging evidence of translational control by AU-rich element-binding proteins. *Front Genet* 2019;10:332.
 61. Guo J, Qu H, Shan T, Chen Y, Chen Y, et al. Tristetraprolin overexpression in gastric cancer cells suppresses PD-L1 expression and inhibits tumor progression by enhancing antitumor immunity. *Mol Cells* 2018;41:653-64.
 62. Deng K, Wang H, Shan T, Chen Y, Zhou H, et al. Tristetraprolin inhibits gastric cancer progression through suppression of IL-33. *Sci Rep* 2016;6:24505.
 63. Sung WW, Chu YC, Chen PR, Liao MH, Lee JW. Positive regulation of HIF-1A expression by EBV oncoprotein LMP1 in nasopharyngeal carcinoma cells. *Cancer Lett* 2016;382:21-31.
 64. Wang M, Yu F, Wu W, Wang Y, Ding H, et al. Epstein-Barr virus-encoded microRNAs as regulators in host immune responses. *Int J Biol Sci* 2018;14:565-76.
 65. Hooykaas MJ, Kruse E, Wiertz EJ, Lebbink RJ. Comprehensive profiling of functional Epstein-Barr virus miRNA expression in human cell lines. *BMC Genomics* 2016;17:644.
 66. Qiu J, Smith P, Leahy L, Thorley-Lawson DA. The Epstein-Barr virus encoded BART miRNAs potentiate tumor growth in vivo. *PLoS Pathog* 2015;11:e1004561.
 67. Yang YC, Liem A, Lambert PF, Sugden B. Dissecting the regulation of EBV's BART miRNAs in carcinomas. *Virology* 2017;505:148-54.
 68. Zhang J, Huang T, Zhou Y, Cheng ASL, Yu J, et al. The oncogenic role of Epstein-Barr virus-encoded microRNAs in Epstein-Barr virus-associated gastric carcinoma. *J Cell Mol Med* 2018;22:38-45.
 69. Tsai CY, Liu YY, Liu KH, Hsu JT, Chen TC, et al. Comprehensive profiling of virus microRNAs of Epstein-Barr virus-associated gastric carcinoma: highlighting the interactions of ebv-Bart9 and host tumor cells. *J Gastroenterol Hepatol* 2017;32:82-91.
 70. Cristino AS, Nourse J, West RA, Sabdia MB, Law SC, et al. EBV microRNA-BHRF1-2-5p targets the 3'UTR of immune checkpoint ligands PD-L1 and PD-L2. *Blood* 2019;134:2261-70.
 71. Skalsky RL, Corcoran DL, Gottwein E, Frank CL, Kang D, et al. The viral and cellular microRNA targetome in lymphoblastoid cell lines. *PLoS Pathog* 2012;8:e1002484.
 72. Vlachos IS, Paraskevopoulou MD, Karagkouni D, Georgakilas G, Vergoulis T, et al. DIANA-TarBase v7.0: indexing more than half a million experimentally supported miRNA:mRNA interactions. *Nucleic Acids Res* 2015;43:D153-9.

73. Yoon CJ, Chang MS, Kim DH, Kim W, Koo BK, et al. Epstein-Barr virus-encoded miR-BART5-5p upregulates PD-L1 through PIAS3/pSTAT3 modulation, worsening clinical outcomes of PD-L1-positive gastric carcinomas. *Gastric Cancer* 2020; epub ahead of print [PMID: 32206940 doi: 10.1007/s10120-020-01059-3]
74. Cerezo M, Guemiri R, Druillennec S, Girault I, Malka-Mahieu H, et al. Translational control of tumor immune escape via the eIF4F-STAT1-PD-L1 axis in melanoma. *Nat Med* 2018;24:1877-86.
75. Jiang X, Zhou J, Giobbie-Hurder A, Wargo J, Hodi FS. The activation of MAPK in melanoma cells resistant to BRAF inhibition promotes PD-L1 expression that is reversible by MEK and PI3K inhibition. *Clin Cancer Res* 2013;19:598-609.
76. Coelho MA, de Carné Trécesson S, Rana S, Zecchin D, Moore C, et al. Oncogenic RAS signaling promotes tumor immunoresistance by stabilizing PD-L1 mRNA. *Immunity* 2017;47:1083-99.e6.
77. Seo AN, Kang BW, Bae HI, Kwon OK, Park KB, et al. Exon 9 mutation of PIK3CA associated with poor survival in patients with Epstein-Barr virus-associated gastric cancer. *Anticancer Res* 2019;39:2145-54.
78. Song M, Chen D, Lu B, Wang C, Zhang J, et al. PTEN loss increases PD-L1 protein expression and affects the correlation between PD-L1 expression and clinical parameters in colorectal cancer. *PLoS One* 2013;8:e65821.
79. Parsa AT, Waldron JS, Panner A, Crane CA, Parney IF, et al. Loss of tumor suppressor PTEN function increases B7-H1 expression and immunoresistance in glioma. *Nat Med* 2007;13:84-8.
80. Kim YB, Ahn JM, Bae WJ, Sung CO, Lee D. Functional loss of ARID1A is tightly associated with high PD-L1 expression in gastric cancer. *Int J Cancer* 2019;145:916-26.
81. Menyhart O, Pongor LS, Györfy B. Mutations defining patient cohorts with elevated PD-L1 expression in gastric cancer. *Front Pharmacol* 2018;9:1522.
82. Mimura K, Teh JL, Okayama H, Shiraishi K, Kua LF, et al. PD-L1 expression is mainly regulated by interferon gamma associated with JAK-STAT pathway in gastric cancer. *Cancer Sci* 2018;109:43-53.
83. Zhang Y, Liu W, Zhang W, Wang W, Song Y, et al. Constitutive activation of the canonical NF- κ B signaling pathway in EBV-associated gastric carcinoma. *Virology* 2019;532:1-10.
84. Shi Q, Zhang Y, Liu W, Xiao H, Qi Y, et al. Latent membrane protein 2A inhibits expression level of Smad2 through regulating miR-155-5p in EBV-associated gastric cancer cell lines. *J Med Virol* 2020;92:96-106.
85. Iwakiri D, Minamitani T, Samanta M. Epstein-Barr virus latent membrane protein 2A contributes to anoikis resistance through ERK activation. *J Virol* 2013;87:8227-34.
86. Qi YF, Liu M, Zhang Y, Liu W, Xiao H, et al. EBV down-regulates COX-2 expression via TRAF2 and ERK signal pathway in EBV-associated gastric cancer. *Virus Res* 2019;272:197735.
87. Li H, Xia JQ, Zhu FS, Xi ZH, Pan CY, et al. LPS promotes the expression of PD-L1 in gastric cancer cells through NF- κ B activation. *J Cell Biochem* 2018;119:9997-10004.
88. Wang X, Wu WKK, Gao J, Li Z, Dong B, et al. Autophagy inhibition enhances PD-L1 expression in gastric cancer. *J Exp Clin Cancer Res* 2019;38:140.
89. Xu D, Li J, Li RY, Lan T, Xiao C, et al. PD-L1 Expression is regulated by NF- κ B during EMT signaling in gastric carcinoma. *Oncotargets Ther* 2019;12:10099-105.
90. Hino R, Uozaki H, Murakami N, Ushiku T, Shinozaki A, et al. Activation of DNA methyltransferase 1 by EBV latent membrane protein 2A leads to promoter hypermethylation of PTEN gene in gastric carcinoma. *Cancer Res* 2009;69:2766-74.
91. Cai L, Ye Y, Jiang Q, Chen Y, Lyu X, et al. Epstein-Barr virus-encoded microRNA BART1 induces tumour metastasis by regulating PTEN-dependent pathways in nasopharyngeal carcinoma. *Nat Commun* 2015;6:7353.
92. Cai LM, Lyu XM, Luo WR, Cui XF, Ye YF, et al. EBV-miR-BART7-3p promotes the EMT and metastasis of nasopharyngeal carcinoma cells by suppressing the tumor suppressor PTEN. *Oncogene* 2015;34:2156-66.
93. Moon JW, Kong SK, Kim BS, Kim HJ, Lim H, et al. IFN γ induces PD-L1 overexpression by JAK2/STAT1/IRF-1 signaling in EBV-positive gastric carcinoma. *Sci Rep* 2017;7:17810.
94. Kawazoe A, Kuwata T, Kuboki Y, Shitara K, Nagatsuma AK, et al. Clinicopathological features of programmed death ligand 1 expression with tumor-infiltrating lymphocyte, mismatch repair, and Epstein-Barr virus status in a large cohort of gastric cancer patients. *Gastric Cancer* 2017;20:407-15.
95. Li Z, Lai Y, Sun L, Zhang X, Liu R, et al. PD-L1 expression is associated with massive lymphocyte infiltration and histology in gastric cancer. *Hum Pathol* 2016;55:182-9.
96. Thompson ED, Zahurak M, Murphy A, Cornish T, Cuka N, et al. Patterns of PD-L1 expression and CD8 T cell infiltration in gastric adenocarcinomas and associated immune stroma. *Gut* 2017;66:794-801.
97. Wallach D, Varfolomeev EE, Malinin NL, Goltsev YV, Kovalenko AV, et al. Tumor necrosis factor receptor and Fas signaling mechanisms. *Annu Rev Immunol* 1999;17:331-67.
98. Billiau A. Interferon- γ : biology and role in pathogenesis. *Advances in Immunology* Volume 62. Elsevier; 1996. pp. 61-130.
99. Jiang X, Wang J, Deng X, Xiong F, Ge J, et al. Role of the tumor microenvironment in PD-L1/PD-1-mediated tumor immune escape. *Mol Cancer* 2019;18:10.
100. Iwai Y, Ishida M, Tanaka Y, Okazaki T, Honjo T, et al. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci U S A* 2002;99:12293-7.
101. Lee SJ, Jang BC, Lee SW, Yang YI, Suh SI, et al. Interferon regulatory factor-1 is prerequisite to the constitutive expression and IFN- γ -induced upregulation of B7-H1 (CD274). *FEBS Lett* 2006;580:755-62.
102. Bellucci R, Martin A, Bommarito D, Wang K, Hansen SH, et al. Interferon- γ -induced activation of JAK1 and JAK2 suppresses tumor cell

- susceptibility to NK cells through upregulation of PD-L1 expression. *Oncoimmunology* 2015;4:e1008824.
103. Ohtani H, Jin Z, Takegawa S, Nakayama T, Yoshie O. Abundant expression of CXCL9 (MIG) by stromal cells that include dendritic cells and accumulation of CXCR3+ T cells in lymphocyte-rich gastric carcinoma. *J Pathol* 2009;217:21-31.
 104. Sasaki S, Nishikawa J, Sakai K, Iizasa H, Yoshiyama H, et al. EBV-associated gastric cancer evades T-cell immunity by PD-1/PD-L1 interactions. *Gastric Cancer* 2019;22:486-96.
 105. Fang W, Zhang J, Hong S, Zhan J, Chen N, et al. EBV-driven LMP1 and IFN- γ up-regulate PD-L1 in nasopharyngeal carcinoma: Implications for oncotargeted therapy. *Oncotarget* 2014;5:12189-202.
 106. Su S, Zou Z, Chen F, Ding N, Du J, et al. CRISPR-Cas9-mediated disruption of PD-1 on human T cells for adoptive cellular therapies of EBV positive gastric cancer. *OncoImmunology* 2016;6:e1249558.
 107. Nakayama A, Abe H, Kunita A, Saito R, Kanda T, et al. Viral loads correlate with upregulation of PD-L1 and worse patient prognosis in Epstein-Barr Virus-associated gastric carcinoma. *PLoS One* 2019;14:e0211358.
 108. Iwakiri D, Zhou L, Samanta M, Matsumoto M, Ebihara T, et al. Epstein-Barr virus (EBV)-encoded small RNA is released from EBV-infected cells and activates signaling from Toll-like receptor 3. *J Exp Med* 2009;206:2091-9.
 109. Lemery S, Keegan P, Pazdur R. First FDA approval agnostic of cancer site - when a biomarker defines the indication. *N Engl J Med* 2017;377:1409-12.