

- bipartite nuclear localization signals in a drug-resistant human lung cancer cell line. *Cancer Res* 1995;55:2129-34.
55. Ritke MK, Yalowich JC. Altered gene expression in human leukemia K562 cells selected for resistance to etoposide. *Biochem Pharmacol* 1993;46:2007-20.
 56. Ritke MK, Roberts D, Allan WP, Raymond J, Bergoltz VV, et al. Altered stability of etoposide-induced topoisomerase II-DNA complexes in resistant human leukemia K562 cells. *Br J Cancer* 1994;69:687-97.
 57. Zwelling LA, Hinds M, Chan D, Mayes J, Sie KL, et al. Characterization of an amsacrine-resistant line of human leukemia cells. Evidence for a drug-resistant form of topoisomerase II. *J Biol Chem* 1989;264:16411-20.
 58. Christie M, Chang CW, Róna G, Smith KM, Stewart AG, et al. Structural biology and regulation of protein import into the nucleus. *J Mol Biol* 2016;428:2060-90.
 59. Gardiner LP, Roper DI, Hammonds TR, Maxwell A. The N-terminal domain of human topoisomerase II alpha is a DNA-dependent ATPase. *Biochemistry* 1998;37:16997-7004.
 60. Campbell S, Maxwell A. The ATP-operated clamp of human DNA topoisomerase II alpha: hyperstimulation of ATPase by "piggy-back" binding. *J Mol Biol* 2002;320:171-88.
 61. Hu T, Sage H, Hsieh TS. ATPase domain of eukaryotic DNA topoisomerase II. Inhibition of ATPase activity by the anti-cancer drug bisdioxopiperazine and ATP/ADP-induced dimerization. *J Biol Chem* 2002;277:5944-51.
 62. Austin CA, Lee KC, Swan RL, Khazeem MM, Manville CM, et al. TOP2B: the first thirty years. *Int J Mol Sci* 2018;19. E2765.
 63. Cowell IG, Austin CA. Do transcription factories and TOP2B provide a recipe for chromosome translocations in therapy-related leukemia? *Cell Cycle* 2012;11:3143-4.
 64. Cowell IG, Sondka Z, Smith K, Lee KC, Manville CM, et al. Model for MLL translocations in therapy-related leukemia involving topoisomerase II β -mediated DNA strand breaks and gene proximity. *Proc Natl Acad Sci U S A* 2012;109:8989-94.