

## Supplementary Materials

**Effects of early life exposure to di(2-ethylhexyl) phthalate on jejunal morphology, sucrase activity, and colonic microbiota composition in young pigs**

**Marcia H. Monaco<sup>1</sup>, Sierra Coyne<sup>1</sup>, Mei Wang<sup>1</sup>, Jodi A. Flaws<sup>2,3</sup>, Joseph M. Irudayaraj<sup>3,4</sup>, Isaac Cann<sup>3,5,6</sup>, Sharon M. Donovan<sup>1,3</sup>**

<sup>1</sup>Department of Food Science and Human Nutrition, College of Agricultural, Consumer and Environmental Sciences (ACES), University of Illinois, Urbana, IL 60801, USA.

<sup>2</sup>Department of Comparative Biosciences, College of Veterinary Medicine, University of Illinois, Urbana, IL 60801, USA.

<sup>3</sup>Institute for Genomic Biology, University of Illinois, Urbana, IL 60801, USA.

<sup>4</sup>Department of Bioengineering, Grainger College of Engineering, University of Illinois, Urbana, IL 60801, USA.

<sup>5</sup>Department of Animal Sciences, College of ACES, University of Illinois, Urbana, IL 60801, USA.

<sup>6</sup>Department of Microbiology, University of Illinois, Urbana, IL 60801, USA.

**Correspondence to:** Dr. Sharon M. Donovan, Department of Food Science and Human Nutrition, College of Agricultural, Consumer and Environmental Sciences (ACES), University of Illinois, 339 Bevier Hall, 905 South Goodwin Ave, Urbana, IL 61801, USA.  
E-mail: [sdonovan@illinois.edu](mailto:sdonovan@illinois.edu)

**Supplementary Table 1. Average daily weight gain (ADG) over various postnatal time periods in piglets supplemented orally with 0 (CON), 20 (DEHP20), or 200 (DEHP200) mg DEHP/kg BW/day**

PND	Treatment groups			<i>P</i> value
	CON	DEHP20	DEHP200	
2-9	155 ± 4.9	152 ± 7.6	150 ± 7.7	0.87
9-16	324 ± 8.9	324 ± 10	293 ± 10	0.07
16-23	469 ± 8.4	479 ± 8.6	484 ± 10	0.48

Data are expressed as mean ± SEM. *N* = 8 per group. ADG was calculated by dividing the total BW gained (g) within each time period by the number of days in the period.

PND: Postnatal day.